PE Conjugated Anti-Mouse CD3 Recombinant Antibody [PSH04-98] - Rabbit IgG (Chimeric)

HA720200F

Product Type: Recombinant Chimeric Antibody IgG, primary antibodies

Species reactivity: Mouse **Applications:** FC

Molecular Wt: Predicted band size: 21 kDa

Clone number: PSH04-98

Description: CD3 (cluster of differentiation 3) is a protein complex and T cell co-receptor that is involved

in activating both the cytotoxic T cell (CD8+ naive T cells) and T helper cells (CD4+ naive T cells). It is composed of four distinct chains. In mammals, the complex contains a CD3 γ chain, a CD3 γ chain, and two CD3 γ chains. These chains associate with the T-cell receptor (TCR) and the CD3-zeta (γ -chain) to generate an activation signal in T lymphocytes. The

TCR, CD3-zeta, and the other CD3 molecules together constitute the TCR complex.

Conjugate: PE

Immunogen: Recombinant protein.

Positive control: Mouse spleen.

Subcellular location: Cell membrane.

Database links: SwissProt: P22646 Mouse

Recommended Dilutions:


FC 5 μl per million cells in 100 μl staining volume or 5 μl per 100 μl of whole blood.

Storage Buffer: Supplied in phosphate-buffered solution, pH 7.2, containing 0.2% ProClean 950 and BSA.

Storage Instruction: Store at 2° to 8° . Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.



Service mail:support@huabio.cn



Images




Fig1: Flow cytometric analysis of mouse spleen labeling Mouse CD3.

Cells were washed twice with cold PBS and resuspend. Then stained with the primary antibody (HA720200F, $1\mu g/mL$) (red). Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- 1. LeFevre JD et al. Anti-CD3 monoclonal antibodies for the prevention and treatment of type 1 diabetes: A literature review. Am J Health Syst Pharm. 2022 Nov
- 2. Chen Y et al. Cholesterol inhibits TCR signaling by directly restricting TCR-CD3 core tunnel motility. Mol Cell. 2022

 Apr