APC Conjugated Anti-Mouse CD3 Recombinant Antibody [PSH04-98] - Rabbit IgG (Chimeric)

HA720201F

Product Type:	Recombinant Chimeric Antibody IgG, primary antibodies
Species reactivity:	Mouse
Applications:	FC
Molecular Wt:	Predicted band size: 21 kDa
Clone number:	PSH04-98
Description:	CD3 (cluster of differentiation 3) is a protein complex and T cell co-receptor that is involved in activating both the cytotoxic T cell (CD8+ naive T cells) and T helper cells (CD4+ naive T cells). It is composed of four distinct chains. In mammals, the complex contains a CD3γ chain, a CD3 δ chain, and two CD3 ϵ chains. These chains associate with the T-cell receptor (TCR) and the CD3-zeta (ζ -chain) to generate an activation signal in T lymphocytes. The TCR, CD3-zeta, and the other CD3 molecules together constitute the TCR complex.
Conjugate:	APC
lmmunogen:	Recombinant protein.
Positive control:	Mouse spleen.
Subcellular location:	Cell membrane.
Database links:	SwissProt: P22646 Mouse
Recommended Dilutions: FC	5 μI per million cells in 100 μI staining volume or 5 μI per 100 μI of whole blood.
Storage Buffer:	Supplied in phosphate-buffered solution, pH 7.2, containing 0.2% ProClean 950 and BSA.
Storage Instruction:	Store at 2 $^\circ\!{\rm C}$ to 8 $^\circ\!{\rm C}$. Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

45 Service mail:support@huabio.cn



11.

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

```
HA720201F - Page 2
```

Images




Fig1: Flow cytometric analysis of mouse spleen labeling Mouse CD3.

Cells were washed twice with cold PBS and resuspend. Then stained with the primary antibody (HA720201F, $1\mu g/mL$) (red). Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- 1. LeFevre JD et al. Anti-CD3 monoclonal antibodies for the prevention and treatment of type 1 diabetes: A literature review. Am J Health Syst Pharm. 2022 Nov
- Chen Y et al. Cholesterol inhibits TCR signaling by directly restricting TCR-CD3 core tunnel motility. Mol Cell. 2022 Apr

Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation