

Anti-Caspase-3 p12 Antibody [JE07-51]

HA722235



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P
Molecular Wt:	Predicted band size: 32/12 kDa
Clone number:	JE07-51

Description: Caspase-3 is a caspase protein that interacts with caspase-8 and caspase-9. It is encoded by the CASP3 gene. CASP3 orthologs have been identified in numerous mammals for which complete genome data are available. Unique orthologs are also present in birds, lizards, lissamphibians, and teleosts. Caspase-3 shares many of the typical characteristics common to all currently-known caspases. For example, its active site contains a cysteine residue (Cys-163) and histidine residue (His-121) that stabilize the peptide bond cleavage of a protein sequence to the carboxy-terminal side of an aspartic acid when it is part of a particular 4-amino acid sequence. This specificity allows caspases to be incredibly selective, with a 20,000-fold preference for aspartic acid over glutamic acid. A key feature of caspases in the cell is that they are present as zymogens, termed procaspases, which are inactive until a biochemical change causes their activation. Each procaspase has an N-terminal large subunit of about 20 kDa followed by a smaller subunit of about 10 kDa, called p20 and p10, respectively.

Immunogen: Recombinant protein within Human Caspase-3 p12 aa 128-277 / 277.

Positive control: HeLa cell lysate, HeLa treated with 1 μ M staurosporine for 4 hours cell lysate, NIH/3T3 cell lysate, NIH/3T3 treated with 1 μ M staurosporine for 3 hours cell lysate, C6 cell lysate, C6 treated with 1 μ M staurosporine for 4 hours cell lysate, human breast cancer tissue, human tonsil tissue, mouse liver tissue, mouse kidney tissue, rat kidney tissue.

Subcellular location: Cytoplasm.

Database links: SwissProt: P42574 Human | P70677 Mouse | P55213 Rat

Recommended Dilutions:

WB	1:1,000
IHC-P	1:200-1:1,000

Storage Buffer: 1*TBS (pH7.4), 0.05% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of Caspase-3 p12 on different lysates with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 1 μ M staurosporine for 4 hours cell lysate

Lane 3: NIH/3T3 cell lysate

Lane 4: NIH/3T3 treated with 1 μ M staurosporine for 3 hours cell lysate

Lane 5: C6 cell lysate

Lane 6: C6 treated with 1 μ M staurosporine for 4 hours cell lysate

Lysates/proteins at 20 μ g/Lane.

Predicted band size: 32/12 kDa


Observed band size: 32/12 kDa

Exposure time: 3 minutes; ECL: K1802;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA722235) at 1/1,000 dilution was used in 5% NFDm/TBST at 4 $^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of Caspase-3 p12 on different lysates with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/2,000 dilution.



Lane 1: HAP1-parental cell lysate (10 μ g/Lane)

Lane 2: HAP1-Caspase-3 p12 KD cell lysate (10 μ g/Lane)

Predicted band size: 32 kDa

Observed band size: 32 kDa

Exposure time: 40 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA722235) at 1/2,000 dilution was used in K1803 at 4 $^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig3: Immunohistochemical analysis of paraffin-embedded human breast cancer tissue with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA722235) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig4: Immunohistochemical analysis of paraffin-embedded human tonsil tissue with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA722235) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig5: Immunohistochemical analysis of paraffin-embedded mouse liver tissue with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA722235) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig6: Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA722235) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig7: Immunohistochemical analysis of paraffin-embedded rat kidney tissue with Rabbit anti-Caspase-3 p12 antibody (HA722235) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA722235) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Dong L et al. Echinacoside Induces Apoptosis in Human SW480 Colorectal Cancer Cells by Induction of Oxidative DNA Damages. *Int J Mol Sci* 16:14655-68 (2015).
2. Nilsonne G et al. Phenotype-dependent apoptosis signalling in mesothelioma cells after selenite exposure. *J Exp Clin Cancer Res* 28:92 (2009).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn