

Biotin Conjugated Anti-Human PD-L2 Antibody [PSH09-82] - Detector

HA723148B



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	ELISA(Det), ELISA
Clone number:	PSH09-82

Description: Involved in negative regulation of activated T cell proliferation; negative regulation of interferon-gamma production; and negative regulation of interleukin-10 production. Predicted to be located in plasma membrane. Predicted to be active in external side of plasma membrane. Biomarker of pulmonary tuberculosis. Involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.

Conjugate: Biotin-conjugated

Immunogen: Recombinant protein within Human PD-L2 Protein aa 20-220 (HA210599)

Positive control: Recombinant Human PD-L2 Protein (HA210599).

Subcellular location: Cell membrane, Secreted.

Database links: SwissProt: Q9BQ51 Human

Recommended Dilutions:

ELISA(Det) Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PSH09-80] to Human PD-L2 antibody (Capture) (HA723145) or Rabbit monoclonal [PSH09-81] to Human PD-L2 antibody (Capture) (HA723146) and Recombinant Human PD-L2 protein (HA210599) as the standard. The reference range value is 31.25-4,000 pg/ml.

ELISA Use at an assay dependent concentration.

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% ProClin300.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation


Images

Fig1: Sandwich ELISA analysis of Human PD-L2 matched pair antibodies




Elisa assay was performed by coating wells of a 96-well plate with 100 μ l per well of capture antibody (HA723145) diluted in carbonate/bicarbonate buffer, at a concentration of 2 μ g/ml overnight at 4 $^{\circ}$ C. Wells of the plate were washed, blocked with 150 μ l 0.05% tween-20 1% BSA blocking buffer, and incubated with serial diluted Recombinant Human Human PD-L2 (HA210599) starting from 4000 pg/ml to 0 pg/ml and detect antibody (HA723148B, 0.2 μ g/ml) for 1 hour at 30 $^{\circ}$ C with shaking. Then the plate was washed and incubated with 100 μ l per well of SA-HRP for 0.5 hour at 30 $^{\circ}$ C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Fig2: Sandwich ELISA analysis of Human PD-L2 matched pair antibodies



Elisa assay was performed by coating wells of a 96-well plate with 100 μ l per well of capture antibody (HA723146) diluted in carbonate/bicarbonate buffer, at a concentration of 2 μ g/ml overnight at 4 $^{\circ}$ C. Wells of the plate were washed, blocked with 150 μ l 0.05% tween-20 1% BSA blocking buffer, and incubated with serial diluted Recombinant Human Human PD-L2 (HA210599) starting from 4000 pg/ml to 0 pg/ml and detect antibody (HA723148B, 0.2 μ g/ml) for 1 hour at 30 $^{\circ}$ C with shaking. Then the plate was washed and incubated with 100 μ l per well of SA-HRP for 0.5 hour at 30 $^{\circ}$ C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Fig3: Interpolated concentrations of native PD-L2 in human urine, human serum samples and Jurkat cell culture supernatant.



The concentrations of PD-L2 were measured in duplicate, interpolated from the PD-L2 standard curve and corrected for sample dilution. Undiluted samples are human urine 100%, human serum 40%, and Jurkat cell culture supernatant 100%. The interpolated dilution factor corrected values are plotted (mean $\pm$ SD, n=2). The mean PD-L2 concentration was determined to be 4,248 pg/ml in human urine, 10,104 pg/ml in human serum and undetectable in Jurkat cell culture supernatant.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation




Fig4: Interpolated concentrations of native PD-L2 in human urine, human serum samples and Jurkat cell culture supernatant.

The concentrations of PD-L2 were interpolated from the PD-L2 standard curve and corrected for sample dilution. Undiluted samples are human urine 100%, human serum 40%, and Jurkat cell culture supernatant 100%. The mean PD-L2 concentration was determined to be 4,983 pg/ml in human urine, 14,438 pg/ml in human serum and undetectable in Jurkat cell culture supernatant.




Fig5: Interpolated concentrations of spiked PD-L2 in human cell culture media samples.

The concentrations of PD-L2 were measured in duplicates, interpolated from the PD-L2 standard curves and corrected for sample dilution. Undiluted samples are as follows: cell culture media 50%. The interpolated dilution factor corrected values are plotted (mean $\pm$ SD, n=2).




Fig6: Interpolated concentrations of spiked PD-L2 in human cell culture media samples.

The concentrations of PD-L2 were measured in duplicates, interpolated from the PD-L2 standard curves and corrected for sample dilution. Undiluted samples are as follows: cell culture media 50%. The interpolated dilution factor corrected values are plotted (mean $\pm$ SD, n=2).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Tseng S.-Y., Otsuji M., Gorski K., Huang X., Slansky J.E., Pai S.I., Shalabi A., Shin T., Pardoll D.M., Tsuchiya H. B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells. J. Exp. Med. 193:839-846 (2001)
2. Latchman Y., Wood C.R., Chernova T., Chaudhary D., Borde M., Chernova I., Iwai Y., Long A.J., Brown J.A., Freeman G.J. PD-L2 is a second ligand for PD-1 and inhibits T cell activation. Nat. Immunol. 2:261-268 (2001)

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation