

# HRP Conjugated Anti-Human ICAM1 Antibody [PSH10-16] - Detector

## HA723474H



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human
<b>Applications:</b>	ELISA(Det), ELISA
<b>Clone number:</b>	PSH10-16

**Description:** ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation. Acts as a receptor for major receptor group rhinovirus A-B capsid proteins. Acts as a receptor for Coxsackievirus A21 capsid proteins. Upon Kaposi's sarcoma-associated herpesvirus/HHV-8 infection, is degraded by viral E3 ubiquitin ligase MIR2, presumably to prevent lysis of infected cells by cytotoxic T-lymphocytes and NK cell. Interacts with MUC1 and promotes cell aggregation in epithelial cells. Interacts with ARHGEF26/SGEF. Interacts (on T cell side) with CD81, CD247 and CD9 at immunological synapses between antigen-presenting cells and T cells.

**Conjugate:** HRP-conjugated

**Immunogen:** Recombinant protein within Human ICAM1 aa 28-480 (HA210857).

**Positive control:** Recombinant Human ICAM1 protein (HA210857).

**Subcellular location:** Membrane.

**Database links:** SwissProt: P05362 Human

### Recommended Dilutions:

**ELISA(Det)** Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PSH10-15] to Human ICAM1 antibody (Capture) (HA723184) and Recombinant Human ICAM1 protein (HA210857) as the standard.

**ELISA** Use at an assay dependent concentration.

**Storage Buffer:** PBS (pH7.4), 0.1% BSA, 40% Glycerol.

**Storage Instruction:** Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

## Images

**Fig1:** Sandwich ELISA analysis of Human ICAM1 matched pair antibodies

Elisa assay was performed by coating wells of a 96-well plate with 100  $\mu$ l per well of capture antibody (HA723184) diluted in carbonate/bicarbonate buffer, at a concentration of 5ug/ml overnight at 4°C. Wells of the plate were washed, blocked with 150  $\mu$ l 0.05% tween-20 1% BSA blocking buffer, and incubated with serial diluted Recombinant Human ICAM1 protein (HA210857) starting from 4000 pg/ml to 0 pg/ml and detect antibody (HA723474H, 0.2  $\mu$ g/ml) for 1 hour at 30°C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

## Background References

- Hayashi T., Takahashi T., Motoya S., Ishida T., Itoh F., Adachi M., Hinoda Y., Imai K. MUC1 mucin core protein binds to the domain 1 of ICAM-1. Digestion 63 Suppl. 1:87-92 (2001)
- van Buul J.D., Allingham M.J., Samson T., Meller J., Boulter E., Garcia-Mata R., Burridge K. RhoG regulates endothelial apical cup assembly downstream from ICAM1 engagement and is involved in leukocyte trans-endothelial migration. J. Cell Biol. 178:1279-1293 (2007)

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation