

Anti-Phospho-PKC alpha/beta II (T638/T641) Antibody [PSH16-27]

HA723825



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IHC-P, IF-Cell, FC, Dot Blot
Molecular Wt:	Predicted band size: 77 kDa
Clone number:	PSH16-27

Description: Members of the protein kinase C (PKC) family play a key regulatory role in a variety of cellular functions including cell growth and differentiation, gene expression, hormone secretion and membrane function. PKCs were originally identified as serine/threonine protein kinases whose activity was dependent on calcium and phospholipids. Diacylglycerols (DAG) and tumor-promoting phorbol esters bind to and activate PKC. PKCs can be subdivided into many different isoforms (α , β I, β II, γ , δ , ϵ , ζ , η , θ , λ /I, μ and ν). Patterns of expression for each PKC isoform differ among tissues and PKC family members exhibit clear differences in their cofactor dependencies. For instance, the kinase activities of PKC δ and ϵ are independent of Ca^{2+} . On the other hand, most of the other PKC members possess phorbol ester-binding activities and kinase activities.

Immunogen: Synthetic phospho-peptide corresponding to residues surrounding Thr638 of Human PKC alpha aa 622-667 / 672.

Positive control: HEK-293 cell lysate, HeLa cell lysate, C6 cell lysate, Mouse brain tissue lysate, Rat brain tissue lysate, NIH/3T3 cell lysate, NIH/3T3 starved overnight then treated with 200nM TPA for 4 hours cell lysate, human breast cancer tissue, HeLa, NIH/3T3, C6.

Subcellular location: Cytoplasm, Cell membrane, Mitochondrion membrane, Nucleus.

Database links: SwissProt: P17252 Human | P05771-2 Human | P20444 Mouse | P68404-2 Mouse | P05696 Rat | P68403-2 Rat

Recommended Dilutions:

WB	1:25,000
IHC-P	1:200
IF-Cell	1:100
FC	1:1,000
Dot Blot	1:5,000

Storage Buffer: 1*PBS (pH7.4), 0.1% BSA, 40% Glycerol, 0.2% Proclean 950.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of Phospho-PKC alpha/beta II (T638/T641) on different lysates with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/25,000 dilution.

Lane 1: HEK-293 cell lysate

Lane 2: HeLa cell lysate

Lane 3: C6 cell lysate

Lane 4: Mouse brain tissue lysate

Lane 5: Rat brain tissue lysate

Lane 6: NIH/3T3 cell lysate

Lane 7: NIH/3T3 starved overnight then treated with 200nM TPA for 4 hours cell lysate

Lane 8: NIH/3T3 starved overnight then treated with 200nM TPA for 4 hours cell lysate, then the membrane treated with λ pp for 1 hour

Lysates/proteins at 20 μ g/Lane.

Predicted band size: 77 kDa

Observed band size: 77 kDa

Exposure time: 14 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (HA723825) at 1/25,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.




Fig2: Immunohistochemical analysis of paraffin-embedded human breast cancer tissue untreated / treated with λ pp with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA723825) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn


Fig3: Immunocytochemistry analysis of HeLa cells untreated / treated with λ pp labeling Phospho-PKC alpha/beta II (T638/T641) with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.


Fig4: Immunocytochemistry analysis of NIH/3T3 cells untreated / treated with λ pp labeling Phospho-PKC alpha/beta II (T638/T641) with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.


Fig5: Immunocytochemistry analysis of C6 cells untreated / treated with λ pp labeling Phospho-PKC alpha/beta II (T638/T641) with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.


Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Fig6: Flow cytometric analysis of NIH/3T3 cells untreated (left) / treated with λ pp (right) labeling Phospho-PKC alpha/beta II (T638/T641).



Cells were fixed and permeabilized. Then stained with the primary antibody (HA723825, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 °C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4 °C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

Fig7: Flow cytometric analysis of C6 cells untreated (left) / treated with λ pp (right) labeling Phospho-PKC alpha/beta II (T638/T641).



Cells were fixed and permeabilized. Then stained with the primary antibody (HA723825, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 °C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4 °C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).




Fig8: Dot blot analysis of Phospho-PKC alpha/beta II (T638/T641) on different peptides with Rabbit anti-Phospho-PKC alpha/beta II (T638/T641) antibody (HA723825) at 1/5,000 dilution. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution for 1 hour at room temperature.

Lane 1: Phospho-PKC alpha (T638) peptide (positive)
 Lane 2: Phospho-PKC beta II (T641) peptide (positive)
 Lane 3: Phospho-PKC beta I (T642) peptide (negative)
 Lane 4: Unmodified PKC alpha peptide (negative)

Proteins loading: 100ng, 25ng, 5ng;

Blocking and dilution buffer: 5% NFDM/TBST;
 Exposure time: 3 seconds; ECL: K1801.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Ghashghaeinia M et al. Coronavirus disease 2019 (COVID-19), human erythrocytes and the PKC-alpha/-beta inhibitor chelerythrine -possible therapeutic implication. Cell Cycle. 2020 Dec

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUAABIO
www.huabio.cn