

Anti-Mouse CD30 Antibody [PSH17-54] - BSA and Azide free (Capture)

HA723929



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Mouse
Applications:	ELISA(Cap)
Clone number:	PSH17-54

Description: CD30, also known as TNFRSF8 (TNF receptor superfamily member 8), is a cell membrane protein of the tumor necrosis factor receptor family and a tumor marker for anaplastic large cell lymphoma. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. It is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Immunogen: Recombinant protein within Mouse CD30 aa 19-287 (HA211062).

Positive control: Recombinant Mouse CD30 protein (HA211062).

Subcellular location: Cell membrane.

Database links: SwissProt: Q60846 Mouse

Recommended Dilutions:

ELISA(Cap) Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PSH17-55] to Mouse CD30 antibody (Detector) (HA723930) and Recombinant Mouse CD30 protein (HA211062) as the standard. The reference range value is 31.2-8,000 pg/mL.

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


 华安生物
HUABIO
www.huabio.cn

Images


Fig1: Sandwich ELISA analysis of Mouse CD30 matched pair antibodies

Capture: HA723929, Mouse CD30 Rabbit mAb [PSH17-54]

Detector: HA723930, Mouse CD30 Rabbit mAb [PSH17-55]




Elisa assay was performed by coating wells of a 96-well plate with 100 μ l per well of capture antibody (HA723929) diluted in carbonate/bicarbonate buffer, at a concentration of 2ug/ml overnight at 4°C. Wells of the plate were washed, blocked with 150 μ l 0.05% tween-20 1% BSA blocking buffer, and incubated with serial diluted Recombinant Mouse CD30 protein (HA211062) starting from 8000 pg/ml to 0 pg/ml and detect antibody (HA723930, Biotin, 0.2 μ g/ml) for 1 hour at 30°C with shaking. Then the plate was washed and incubated with 100 μ l per well of SA-HRP for 0.5 hour at 30°C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Fig2: Interpolated concentrations of native CD30 in mouse spleen cell supernatant treated or untreated with IL-2 and Concanavalin A.

Capture: HA723929, Mouse CD30 Rabbit mAb [PSH17-54]

Detector: HA723930, Mouse CD30 Rabbit mAb [PSH17-55]

Interpolated concentration of native CD30 was measured in duplicate at different sample concentrations and interpolated from the CD30 standard curves. The interpolated dilution factor corrected values were plotted (mean \pm SD, n=2). The mean CD30 concentration was determined to be 9,657 pg/mL in treated mouse spleen cell supernatant. There was no detectable signal in untreated mouse spleen cell supernatant.

Fig3: Interpolated concentrations of spiked CD30 in cell culture media samples.

Capture: HA723929, Mouse CD30 Rabbit mAb [PSH17-54]

Detector: HA723930, Mouse CD30 Rabbit mAb [PSH17-55]

The concentrations of CD30 were measured in duplicates, interpolated from the Osteopontin CD30 standard curves and corrected for sample dilution. Diluted samples are as follows: 50% cell culture media with FBS. The interpolated dilution factor corrected values are plotted (mean \pm SD, n=2).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Horwitz S et al. The ECHELON-2 Trial: 5-year results of a randomized, phase III study of brentuximab vedotin with chemotherapy for CD30-positive peripheral T-cell lymphoma. *Ann Oncol.* 2022 Mar
2. Nakashima M et al. CD30 Expression and Its Functions during the Disease Progression of Adult T-Cell Leukemia/Lymphoma. *Int J Mol Sci.* 2023 May

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn