

Anti-Human DPP4 / CD26 Antibody [PSH19-28] - BSA and Azide free (Capture)

HA724060



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	ELISA(Cap)
Clone number:	PSH19-28

Description: Dipeptidyl peptidase-4 (DPP4 or DPPIV), also known as adenosine deaminase complexing protein 2 or CD26 (cluster of differentiation 26) is a protein that, in humans, is encoded by the DPP4 gene. DPP4 is related to FAP, DPP8, and DPP9. The enzyme was discovered in 1966 by Hopsu-Havu and Glenner, and as a result of various studies on chemism, was called dipeptidyl peptidase IV [DP IV]. The protein encoded by the DPP4 gene is an enzyme expressed on the surface of most cell types and is associated with immune regulation, signal transduction, and apoptosis. It is a type II transmembrane glycoprotein, but a soluble form, which lacks the intracellular and transmembrane part, is present in blood plasma and various body fluids. DPP-4 is a serine exopeptidase that cleaves X-proline or X-alanine dipeptides from the N-terminus of polypeptides. Peptide bonds involving the cyclic amino acid proline cannot be cleaved by the majority of proteases and an N-terminal X-proline "shields" various biopeptides.[7] Extracellular proline-specific proteases therefore play an important role in the regulation of these biopeptides. DPP-4 is known to cleave a broad range of substrates including growth factors, chemokines, neuropeptides, and vasoactive peptides. The cleaved substrates lose their biological activity in the majority of cases, but in the case of the chemokine RANTES and neuropeptide Y, DPP-4 mediated cleavage leads to a shift in the receptor subtype binding.

Immunogen:	Recombinant protein within Human DPP4 aa 29-766 (HA211034).
Positive control:	Recombinant Human DPP4 / CD26 protein (HA211034).
Subcellular location:	Secreted, Cell membrane, Apical cell membrane, invadopodium membrane, lamellipodium membrane, Cell junction, Membrane raft.
Database links:	SwissProt: P27487 Human
Recommended Dilutions:	
ELISA(Cap)	Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PSH19-27] to Human DPP4 / CD26 antibody (Detector) (HA724059) and Recombinant Human DPP4 / CD26 protein (HA211034) as the standard. The reference range value is 39.1-10,000 pg/mL.
Storage Buffer:	1*PBS (pH7.4).
Storage Instruction:	Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物
HUABIO
www.huabio.cn

Images

Fig1: Sandwich ELISA analysis of Human DPP4 matched pair antibodies


Capture: HA724060, Human DPP4 / CD26 Rabbit mAb [PSH19-28]

Detector: HA724059, Human DPP4 / CD26 Rabbit mAb [PSH19-27]



Elisa assay was performed by coating wells of a 96-well plate with 100 μ l per well of capture antibody (HA724060) diluted in carbonate/bicarbonate buffer, at a concentration of 5 μ g/mL overnight at 4°C. Wells of the plate were washed, blocked with 150 μ l 0.05% tween-20 1% BSA blocking buffer, and incubated with serial diluted Recombinant Human DPP4 / CD26 protein (HA211034) starting from 10,000 pg/ml to 0 pg/ml and detect antibody (HA724059, Biotin, 0.2 μ g/ml) for 1 hour at 30°C with shaking. Then the plate was washed and incubated with 100 μ l per well of SA-HRP for 0.5 hour at 30°C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Fig2: Interpolated concentrations of native DPP4 in HepG2, MCF-7 culture supernatant and Human urine.



Capture: HA724060, Human DPP4 / CD26 Rabbit mAb [PSH19-28]

Detector: HA724059, Human DPP4 / CD26 Rabbit mAb [PSH19-27]

Interpolated concentration of native DPP4 was measured in duplicate at different sample concentrations and interpolated from the DPP4 standard curves. The interpolated dilution factor corrected values were plotted (mean \pm SD, n=2). The mean DPP4 concentration was determined to be 25,703 pg/ml in Human urine, 5,564 pg/ml in HepG2 cell culture supernatant and undetectable in MCF-7 cell culture supernatant.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig3: Interpolated concentrations of spiked DPP4 in cell culture media samples.

Capture: HA724060, Human DPP4 / CD26 Rabbit mAb [PSH19-28]

Detector: HA724059, Human DPP4 / CD26 Rabbit mAb [PSH19-27]

The concentrations of DPP4 were measured in duplicates, interpolated from the DPP4 standard curves and corrected for sample dilution. Undiluted samples are as follows: cell culture media 50%. The interpolated dilution factor corrected values are plotted (mean +/- SD, n=2).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Love KM et al. DPP4 Activity, Hyperinsulinemia, and Atherosclerosis. J Clin Endocrinol Metab. 2021 May
2. Chen SY et al. DPP4 as a Potential Candidate in Cardiovascular Disease. J Inflamm Res. 2022 Sep

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn