

HRP Conjugated Anti-Human/Monkey Albumin Antibody [PSH20-17] - Detector

HA724128H



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Dog, Cynomolgus monkey
Applications:	ELISA(Det), ELISA
Clone number:	PSH20-17

Description: Serum albumin (ALB), the main protein in plasma, has a very good binding capacity for water, fatty acids, calcium, sodium, bilirubin, hormones, potassium and drugs. The primary function of ALB is to regulate the colloidal osmotic pressure of blood. Albumin is synthesized in the liver as preproalbumin, which has an N-terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to produce the secreted form of albumin. Mutations in the ALB gene may result in familial dysalbuminemic hyperthyroxinemia (FDH), a form of euthyroid hyperthyroxinemia that is due to increased affinity of ALB for T4. FDH is the most common cause of inherited euthyroid hyperthyroxinemia in Caucasian populations.

Conjugate: HRP-conjugated

Immunogen: Recombinant protein within Human Albumin.

Positive control: Human Albumin protein (A3782, sigma).

Subcellular location: Secreted.

Database links: SwissProt: P02768 Human

Recommended Dilutions:

ELISA(Det) Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PSH20-16] to Human/Monkey Albumin antibody (Capture) (HA724127) and Human Albumin protein (A3782, sigma) as the standard. The reference range value is 39.06-5,000 pg/mL.

ELISA Use at an assay dependent concentration.

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn


Images

Fig1: Sandwich ELISA analysis of Human Albumin matched pair antibodies

Capture: HA724127, Human/Monkey Albumin Rabbit mAb [PSH20-16]

Detector: HA724128, Human/Monkey Albumin Rabbit mAb [PSH20-17]

Standard curve of Human Albumin matched pair antibodies



Elisa assay was performed by coating wells of a 96-well plate with 100 μ l per well of capture antibody (HA724127) diluted in carbonate/bicarbonate buffer, at a concentration of 2 μ g/mL overnight at 4 $^{\circ}$ C. Wells of the plate were washed, blocked with 150 μ l 0.05% tween-20 1% BSA blocking buffer, and incubated with serial diluted Human Albumin protein (A3782, sigma) starting from 5,000 pg/ml to 0 pg/ml and detect antibody (HA724128, HRP, 0.05 μ g/ml) for 1 hour at 30 $^{\circ}$ C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Zhai W et al. A1 adenosine receptor attenuates intracerebral hemorrhage-induced secondary brain injury in rats by activating the P38-MAPKAP2-Hsp27 pathway. *Mol Brain* 9:66 (2016).
2. Ma J et al. Pramipexole-Induced Hypothermia Reduces Early Brain Injury via PI3K/AKT/GSK3 pathway in Subarachnoid Hemorrhage rats. *Sci Rep* 6:23817 (2016).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn