

Anti-Phospho-c-Fos (S32) Antibody [PSH21-99] HA724268



Product Type: Recombinant Rabbit monoclonal IgG, primary antibodies

Species reactivity: Human

Applications: WB, IF-Cell

Molecular Wt: Predicted band size: 41 kDa

Clone number: PSH21-99

Description: Protein c-Fos is a proto-oncogene that is the human homolog of the retroviral oncogene *v-fos*. It is encoded in humans by the FOS gene. It was first discovered in rat fibroblasts as the transforming gene of the FBJ MSV (Finkel–Biskis–Jinkins murine osteogenic sarcoma virus). It is a part of a bigger Fos family of transcription factors which includes c-Fos, FosB, Fra-1 and Fra-2. It has been mapped to chromosome region 14q21→q31. c-Fos encodes a 62 kDa protein, which forms heterodimer with c-jun (part of Jun family of transcription factors), resulting in the formation of AP-1 (Activator Protein-1) complex which binds DNA at AP-1 specific sites at the promoter and enhancer regions of target genes and converts extracellular signals into changes of gene expression. It plays an important role in many cellular functions and has been found to be overexpressed in a variety of cancers.

Immunogen: Synthetic phospho-peptide corresponding to residues surrounding Ser32 of Human c-Fos.

Positive control: HeLa starved for 16 hours then treated with 200nM TPA for 4 hours cell lysate, HeLa cells starved for 16 hours then treated with 200nM TPA for 4 hours.

Subcellular location: Nucleus, Endoplasmic reticulum, Cytoplasm, cytosol.

Database links: SwissProt: P01100 Human

Recommended Dilutions:

WB 1:5,000

IF-Cell 1:1,000

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of Phospho-c-Fos (S32) on different lysates with Rabbit anti-Phospho-c-Fos (S32) antibody (HA724268) at 1/5,000 dilution.



Lane 1: HeLa starved for 16 hours cell lysate
 Lane 2: HeLa starved for 16 hours then treated with 200nM TPA for 4 hours cell lysate
 Lane 3: HeLa starved for 16 hours then treated with 200nM TPA for 4 hours cell lysate, then the membrane treated with λpp for 1 hour


Lysates/proteins at 15 µg/Lane.

Predicted band size: 41 kDa
 Observed band size: 55 kDa

Exposure time: 42 seconds; ECL: K1801;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA724268) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Immunocytochemistry analysis of HeLa cells untreated / starved for 16 hours then treated with 200nM TPA for 4 hours labeling Phospho-c-Fos (S32) with Rabbit anti-Phospho-c-Fos (S32) antibody (HA724268) at 1/1,000 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Phospho-c-Fos (S32) antibody (HA724268) at 1/1,000 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Matsuoka K et al. Metabolic rewiring controlled by c-Fos governs cartilage integrity in osteoarthritis. *Ann Rheum Dis.* 2023 Sep
2. Osada N et al. c-FOS is an integral component of the IKZF1 transactivator complex and mediates lenalidomide resistance in multiple myeloma. *Clin Transl Med.* 2023 Aug

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn