

# HRP Conjugated Anti-Human/Monkey VEGFR3 / FLT4 Antibody [PSH22-17] - Detector HA724278H



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Cynomolgus monkey
<b>Applications:</b>	ELISA(Det), ELISA
<b>Clone number:</b>	PSH22-17

**Description:** Fms-related tyrosine kinase 4, also known as FLT4, is a protein which in humans is encoded by the FLT4 gene. This gene encodes a tyrosine kinase receptor for vascular endothelial growth factors C and D. The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA.

**Conjugate:** HRP-conjugated

**Immunogen:** Recombinant protein within Human VEGFR3 / FLT4 aa 25-775 (HA211107).

**Positive control:** Recombinant Human VEGFR3 / FLT4 protein (HA211107).

**Subcellular location:** Cell membrane, Cytoplasm, Nucleus.

**Database links:** SwissProt: P35916 Human

## Recommended Dilutions:

**ELISA(Det)** Use at an assay dependent concentration. Can be paired for Sandwich ELISA with Rabbit monoclonal [PSH22-16] to Human/Monkey VEGFR3 / FLT4 antibody (Capture) (HA724277) and Recombinant Human VEGFR3 / FLT4 protein (HA211107) as the standard. The reference range value is 78.1-10,000 pg/mL.  
**ELISA** Use at an assay dependent concentration.

**Storage Buffer:** 1\*PBS (pH7.4), 0.1% BSA, 40% Glycerol, 0.2% Proclean 950.

**Storage Instruction:** Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn


## Images

**Fig1:** Sandwich ELISA analysis of Human VEGFR3 matched pair antibodies

Capture: HA724277, Human/Monkey VEGFR3 / FLT4 Rabbit mAb [PSH22-16]

Detector: HA724278, Human/Monkey VEGFR3 / FLT4 Rabbit mAb [PSH22-17]

Standard curve of Human VEGFR3 / FLT4 matched pair antibodies



Elisa assay was performed by coating wells of a 96-well plate with 100  $\mu$ l per well of capture antibody (HA724277) diluted in carbonate/bicarbonate buffer, at a concentration of 2  $\mu$ g/mL overnight at 4°C. Wells of the plate were washed, blocked with 150  $\mu$ l 0.2% tween-20 1% BSA blocking buffer, and incubated with serial diluted Recombinant Human VEGFR3 / FLT4 protein (HA211107) starting from 10,000 pg/ml to 0 pg/ml and detect antibody (HA724278, HRP, 0.2  $\mu$ g/ml) for 1 hour at 30°C with shaking. Detection was performed using an Ultra TMB Substrate for 10 minutes at room temperature in the dark. The reaction was stopped with sulfuric acid and absorbances were read on a spectrophotometer at 450 nm.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

## Background References

1. Ma L et al. FLT4/VEGFR3 activates AMPK to coordinate glycometabolic reprogramming with autophagy and inflammasome activation for bacterial elimination. *Autophagy*. 2022 Jun
2. Monaghan RM et al. FLT4 causes developmental disorders of the cardiovascular and lymphovascular systems via pleiotropic molecular mechanisms. *Cardiovasc Res*. 2024 Sep

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn