

Anti-PD-L1 Antibody [PSH22-67]

HA724313



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	IHC-P
Molecular Wt:	Predicted band size: 33 kDa
Clone number:	PSH22-67

Description: PD-L1 (programmed-death ligand 1; CD274), is a transmembrane protein constitutively expressed on a variety of cell types, including antigen presenting cells (dendritic cells and histiocytes) and some non-lymphoid tissues (heart and lung). Binding of PD-L1 to PD-1 (programmed-death 1; CD279) expressed by activated T-cells, inhibits their function, causing negative feedback control of immunological reactions, thus impeding inflammation and autoimmunity. Tumour cells may express PD-L1, which binds to PD-1 allowing cancer cells to evade the attack of T-cells. Blockade of the PD-1/PD-L1 pathway has now shown useful in therapy of multiple cancer types, causing durable tumour regressions in a substantial proportion of otherwise treatment refractory cases of melanoma, and carcinomas of e.g., lung, kidney, and urinary tract. Patients without tumour PD-L1 expression can also derive benefit from blocking agents (studies across multiple cancer types demonstrate a pooled response rate of 48% in patients with PD-L1-positive tumours compared to 15% in PD-L1-negative tumours). Tonsil and placenta can be used as positive and negative tissue controls. However, tonsil is found to be superior to placenta, as tonsil displays a range of PD-L1 expression levels. Tonsil displays the following reaction pattern: No staining reaction in the vast majority of lymphocytes including mantle zone and germinal centre B-cells, no staining reaction in superficial epithelial cells, a weak to moderate, typically punctuated membranous staining reaction of the majority of germinal centre macrophages and finally a moderate to strong staining reaction of the majority of epithelial crypt cells.

Positive control: Human lung carcinoma tissue, human placenta tissue.

Subcellular location: Cell membrane, Early endosome membrane, Recycling endosome membrane, Nucleus.

Database links: SwissProt: Q9NZQ7 Human

Recommended Dilutions:

IHC-P 1:4,000

Storage Buffer: PBS (pH7.4), 0.1% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue with Rabbit anti-PD-L1 antibody (HA724313) at 1/4,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA724313) at 1/4,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig2: Immunohistochemical analysis of paraffin-embedded human placenta tissue with Rabbit anti-PD-L1 antibody (HA724313) at 1/4,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA724313) at 1/4,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Lei Q et al. Resistance Mechanisms of Anti-PD1/PDL1 Therapy in Solid Tumors. *Front Cell Dev Biol.* 2020 Jul
2. Tamene W et al. PDL1 expression on monocytes is associated with plasma cytokines in Tuberculosis and HIV. *PLoS One.* 2021 Oct

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


 华安生物
 HUABIO
 www.huabio.cn