

Anti-Phospho-EGFR (Y992) Antibody [PSH23-63]

HA724388



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	WB, IF-Cell, FC
Molecular Wt:	Predicted band size: 134 kDa
Clone number:	PSH23-63

Description: The epidermal growth factor receptor (EGFR; ErbB-1; HER1 in humans) is a transmembrane protein that is a receptor for members of the epidermal growth factor family (EGF family) of extracellular protein ligands. The epidermal growth factor receptor is a member of the ErbB family of receptors, a subfamily of four closely related receptor tyrosine kinases: EGFR (ErbB-1), HER2/neu (ErbB-2), Her 3 (ErbB-3) and Her 4 (ErbB-4). In many cancer types, mutations affecting EGFR expression or activity could result in cancer. Deficient signaling of the EGFR and other receptor tyrosine kinases in humans is associated with diseases such as Alzheimer's, while over-expression is associated with the development of a wide variety of tumors. Interruption of EGFR signalling, either by blocking EGFR binding sites on the extracellular domain of the receptor or by inhibiting intracellular tyrosine kinase activity, can prevent the growth of EGFR-expressing tumours and improve the patient's condition.

Immunogen: Synthetic phosphopeptide corresponding to residues surrounding Tyr992 of human EGFR.

Positive control: A431 treated with 100ng/mL EGF for 30min cell lysate.

Subcellular location: Cell membrane, Nucleus membrane, Nucleus, Endoplasmic reticulum membrane, Golgi apparatus membrane, Endosome.

Database links: SwissProt: P00533 Human

Recommended Dilutions:

WB	1:2,000
IF-Cell	1:1,000
FC	1:10,000

Storage Buffer: 1*PBS (pH7.4), 0.1% BSA, 40% Glycerol, 0.2% Proclean 950.

Storage Instruction: Shipped at 4°C. Store at +4°C short term (1-2 weeks). It is recommended to aliquot into single-use upon delivery. Store at -20°C long term.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of Phospho-EGFR (Y992) on different lysates with Rabbit anti-Phospho-EGFR (Y992) antibody (HA724388) at 1/2,000 dilution.

Lane 1: A431 (Human epidermoid carcinoma skin squamous cells) cell lysate

Lane 2: A431 treated with 100ng/mL EGF for 30min cell lysate

Lysates/proteins at 20 µg/Lane.

Exposure time: 12 seconds; ECL: K1801

Blocking: 5% NFDm/TBST, 1 hour at room temperature

Primary antibody: HA724388, 1/2,000 in primary antibody dilution buffer (K1803), 2 hours at room temperature

Secondary antibody: Goat anti-Rabbit IgG-HRP (HA1001), 1/50,000 in 5% NFDm/TBST, 1 hour at room temperature


Predicted band size: 134 kDa

Observed band size: 175 kDa

Fig2: Application: Immunocytochemistry (IF-cell)

Species: Human

Sample: A431 (Human epidermoid carcinoma skin squamous cells) treated with EGF(100ng/ml 30min)



Fixation: 4% Paraformaldehyde, 15 minutes at room temperature.

Permeabilization: 0.1% Triton X-100, 15 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 1 hour at room temperature.

Antibody dilution buffer: 1% BSA in PBST.

Primary antibody: HA724388, 1/1,000, overnight at 4°C.

Secondary antibody: Goat Anti-Rabbit IgG (iFluor™ 488, HA1121), 45 minutes at room temperature.

Counterstain: Beta tubulin (HA601187, red), 1/100, overnight at 4°C. The Nuclear counterstain was DAPI (Blue).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Fig3: Application: Flow Cytometry (Intra)

Species: Human

Sample: A431 (Human epidermoid carcinoma skin squamous cells) treated with EGF(100ng/ml 30min)



Fixation: 4% Paraformaldehyde, 15 minutes at room temperature.
 Permeabilization: 1x Permeabilization Buffer (eBioscience), 15 minutes at room temperature.

Blocking: 1% BSA + 10% normal goat serum, 15 minutes at room temperature.

Antibody dilution buffer: 1x PBS.

Primary antibody: HA724388 (1/10,000) (Red) compared with Rabbit IgG Isotype Control (HA722127, Green), 15 minutes at room temperature.

Secondary antibody: Goat Anti-Rabbit IgG (iFluor™ 488, HA1121), 15 minutes at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Nakamura T et al. LRIG1 inhibits STAT3-dependent inflammation to maintain corneal homeostasis. *J Clin Invest* 124:385-97 (2014).
2. Furcht CM et al. Multivariate signaling regulation by SHP2 differentially controls proliferation and therapeutic response in glioma cells. *J Cell Sci* 127:3555-67 (2014).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn