

Anti-Cytokeratin 17 Antibody [SR45-06] - BSA and Azide free

HA750031



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB, IF-Cell, IF-Tissue, IHC-P
Molecular Wt:	Predicted band size: 48 kDa
Clone number:	SR45-06

Description: Cytokeratin 17 is a member of the Cytokeratin subfamily of intermediate filament proteins (IFPs). It is unique in that it is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Cytokeratin 17 contains 432 amino acids and is expressed in the nail bed, hair follicle, sebaceous glands and other epidermal appendages. Cytokeratin 17 functions to regulate cell growth and size through its interactions with the adaptor protein 14-3-3-sigma to mediate protein synthesis. Mutations in the gene encoding for Cytokeratin 17 lead to depressed protein translation and smaller sized skin keratinocytes, corresponding to decreased Akt/mTOR signaling activity. Cytokeratin 17 may be a useful marker for cervical stem cell identification, squamous cell carcinoma of the larynx, respiratory syncytial virus and transitional cell carcinomas of the human urinary tract.

Immunogen: Synthetic peptide within Human Cytokeratin 17 aa 1-50 / 432.

Positive control: A431 cell lysates, Mouse skin tissue lysate, Rat skin tissue lysate, SiHa, human prostate tissue, human skin tissue, human tonsil tissue, mouse prostate tissue, human cervical carcinoma tissue.

Subcellular location: Cytoplasm.

Database links: SwissProt: Q04695 Human | Q9QWL7 Mouse | Q6IFU8 Rat

Recommended Dilutions:

WB	1:1,000-1:2,000
IF-Cell	1:50
IF-Tissue	1:200
IHC-P	1:50-1:1,500

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of Cytokeratin 17 on A431 cell lysates with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/2,000 dilution.

Lysates/proteins at 15 µg/Lane.

Predicted band size: 48 kDa

Observed band size: 48 kDa

Exposure time: 3 minutes;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750031) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of Cytokeratin 17 on different lysates with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,000 dilution.

Lane 1: Mouse skin tissue lysate

Lane 2: Rat skin tissue lysate



Lysates/proteins at 40 µg/Lane.

Predicted band size: 48 kDa

Observed band size: 48 kDa

Exposure time: 3 minutes; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750031) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Fig3: Immunocytochemistry analysis of SiHa cells labeling Cytokeratin 17 with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,000 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,000 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.




Fig4: Immunohistochemical analysis of paraffin-embedded human prostate tissue using anti-Cytokeratin 17 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031, 1/400) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig5: Immunohistochemical analysis of paraffin-embedded human skin tissue with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031) at 1/1,500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig6: Immunohistochemical analysis of paraffin-embedded human skin tissue with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,500 dilution.

The section was not undergone antigen retrieval. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031) at 1/1,500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig7: Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Cytokeratin 17 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig8: Immunohistochemical analysis of paraffin-embedded mouse prostate tissue using anti-Cytokeratin 17 antibody. The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 8.0-8.4) for 20 minutes. The tissues were blocked in 5% BSA for 30 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031, 1/50) for 30 minutes at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig9: Immunohistochemical analysis of paraffin-embedded human cervical carcinoma tissue with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031) at 1/1,500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig10: Immunohistochemical analysis of paraffin-embedded human cervical carcinoma tissue with Rabbit anti-Cytokeratin 17 antibody (HA750031) at 1/1,500 dilution.

The section was not undergone antigen retrieval. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750031) at 1/1,500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig11: Immunofluorescence analysis of paraffin-embedded human skin tissue labeling Cytokeratin 17 (HA750031).

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS. And then probed with the primary antibodies Cytokeratin 17 (HA750031, red) at 1/200 dilution at +4 °C overnight, washed with PBS.

Goat Anti-Rabbit IgG H&L (iFluor™ 594, HA1122) was used as the secondary antibodies at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Doucet YS et al. The touch dome defines an epidermal niche specialized for mechanosensory signaling. *Cell Rep* 3:1759-65 (2013).
2. Johnson EK et al. Identification of new dystroglycan complexes in skeletal muscle. *PLoS One* 8:e73224 (2013).

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
 HUABIO
www.huabio.cn