

## Anti-Phospho-Erk1 (T202)+Erk2 (T185) Antibody [SZ2-4] - BSA and Azide free

# HA750061



|                            |   |
|----------------------------|---|
| <b>Product Type:</b>       | Recombinant Rabbit monoclonal IgG, primary antibodies |
| <b>Species reactivity:</b> | Human, Mouse, Rat                                     |
| <b>Applications:</b>       | WB, IF-Cell, IHC-P, FC, IF-Tissue                     |
| <b>Molecular Wt:</b>       | Predicted band size: 43/41 kDa                        |
| <b>Clone number:</b>       | SZ2-4   |

|                               |   |
|-------------------------------|---|
| <b>Description:</b>           | Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs, such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli including mitogens, growth factors, and cytokines, and research investigators consider it an important target in the diagnosis and treatment of cancer. Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase kinase (MAPKKK or MAP3K), a MAP kinase kinase (MAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family, as well as Mos and Tpl2/COT. MEK1 and MEK2 are the primary MAPKKs in this pathway. MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK and the transcription factor Elk-1. p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known as DUSPs or MKPs, along with MEK inhibitors, such as U0126 and PD98059. |
| <b>Immunogen:</b>             | Synthetic phospho-peptide corresponding to residues surrounding Thr185 of Human Erk2.   |
| <b>Positive control:</b>      | Jurkat treated with 200ng/mL PMA for 35 minutes cell lysate, NIH/3T3 cell lysate, NIH/3T3 treated with 200nM PMA for 30 minutes cell lysate, C6 cell lysate, C6 treated with 200nM PMA for 30 minutes cell lysate, A549, NIH/3T3, MCF-7, human lung carcinoma tissue, human kidney tissue, human gallbladder tissue.  |
| <b>Subcellular location:</b>  | Cytoplasm, Nucleus, Membrane, Cell junction.  |
| <b>Database links:</b>        | SwissProt: P27361 Human   P28482 Human   P63085 Mouse   Q63844 Mouse  |
| <b>Recommended Dilutions:</b> |   |
| WB                            | 1:1,000   |
| IF-Cell                       | 1:100-1:500   |
| IHC-P                         | 1:200-1:1,000   |
| FC                            | 1:50-1:100  |
| IF-Tissue                     | 1:50-1:200  |
| <b>Storage Buffer:</b>        | PBS (pH7.4).  |
| <b>Storage Instruction:</b>   | Store at +4℃ after thawing. Aliquot store at -20℃ or -80℃. Avoid repeated freeze / thaw cycles.   |
| <b>Purity:</b>                | Protein A affinity purified.  |

## Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

## Images



**Fig1:** Western blot analysis of Phospho-Erk1 (T202)+Erk2 (T185) on different lysates with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/1,000 dilution.

Lane 1: Jurkat cell lysate

Lane 2: Jurkat treated with 200ng/mL PMA for 35 minutes cell lysate

Lane 3: NIH/3T3 cell lysate

Lane 4: NIH/3T3 treated with 200nM PMA for 30 minutes cell lysate

Lane 5: C6 cell lysate

Lane 6: C6 treated with 200nM PMA for 30 minutes cell lysate

Lysates/proteins at 10 µg/Lane.

Predicted band size: 41/43 kDa

Observed band size: 41/43 kDa

Exposure time: 3 minutes; ECL: K1802;

4-20% SDS-PAGE gel.


Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750061) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Western blot analysis of Phospho-Erk1(T202)+Erk2(T185) on jurkat cell lysates.

Lane 1: jurkat cells, whole cell lysate, 10ug/lane

Lane 2/3: jurkat cells treated with 200 ng/ml PMA for 35 minutes, whole cell lysate, 10ug/lane

Lane 4: jurkat cells treated with 200 ng/ml PMA for 35 minutes, then treated with 2.8ug/ul lambda-PP for 30 minutes, whole cell lysates, 10ug/lane



All lanes :

Anti-Phospho-Erk1(T202)+Erk2(T185) antibody (HA750061) at 1/500 dilution. Anti-Erk1+Erk2 antibody (ET1601-29) at 1/500 dilution. Anti-GAPDH antibody (ET1601-4) at 1/10,000 dilution. Goat Anti-Rabbit IgG H&L (HRP) (HA1001) at 1/200,000 dilution.

Predicted band size: 41/43 kDa

Observed band size: 41/43 kDa

Blocking and diluting buffer: 5% BSA.

Exposure time: Lan1/2 4 minutes; Lan3/4 3 minutes


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物  
HUABIO  
www.huabio.cn




**Fig3:** ICC staining of Phospho-Erk1 (T202)+Erk2 (T185) in A549 cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody (HA750061, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).




**Fig4:** ICC staining of Phospho-Erk1 (T202)+Erk2 (T185) in NIH/3T3 cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody (HA750061, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).



**Fig5:** ICC staining of Phospho-Erk1 (T202)+Erk2 (T185) in MCF-7 cells (green). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody (HA750061, 1/50) for 1 hour at room temperature, washed with PBS. Alexa Fluor®488 Goat anti-Rabbit IgG was used as the secondary antibody at 1/1,000 dilution. The nuclear counter stain is DAPI (blue).




**Fig6:** Flow cytometric analysis of Phospho-Erk1 (T202)+Erk2 (T185) was done on MCF-7 cells. The cells were fixed, permeabilized and stained with the primary antibody (HA750061, 1/50) (red). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor 488-conjugated Goat anti-Rabbit IgG Secondary antibody at 1/1000 dilution for 30 minutes. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).




**Fig7:** Immunohistochemical analysis of paraffin-embedded human gallbladder tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/100 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/100 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig8:** Immunohistochemical analysis of paraffin-embedded human kidney tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/200 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."



**Fig9:** Immunohistochemical analysis of paraffin-embedded human spleen tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/200 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."






**Fig10:** Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."




**Fig11:** Immunohistochemical analysis of paraffin-embedded mouse spleen tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."




**Fig12:** Immunohistochemical analysis of paraffin-embedded rat kidney tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."



**Fig13:** Immunohistochemical analysis of paraffin-embedded rat spleen tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."



**Fig14:** Immunohistochemical analysis of paraffin-embedded human thyroid cancer tissue with Rabbit anti-Phospho-Erk1 (T202)+Erk2 (T185) antibody (HA750061) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750061) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX."

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Ruess DA et al. HDACi Valproic Acid (VPA) and Suberoylanilide Hydroxamic Acid (SAHA) Delay but Fail to Protect against Warm Hepatic Ischemia-Reperfusion Injury. PLoS One 11:e0161233 (2016).
2. Ahnstedt H et al. U0126 attenuates cerebral vasoconstriction and improves long-term neurologic outcome after stroke in female rats. J Cereb Blood Flow Metab 35:454-60 (2015).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUAABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation