

Anti-TBR1 Antibody [JF10-00] - BSA and Azide free

HA750366



Product Type: Recombinant Rabbit monoclonal IgG, primary antibodies

Species reactivity: Human, Mouse, Rat, Cynomolgus monkey, Pig

Applications: WB, IHC-P, IF-Tissue, IHC-Fr

Molecular Wt: Predicted band size: 74 kDa

Clone number: JF10-00

Description: A novel murine and human gene, TBR-1, encodes a putative transcription factor related to the Brachyrury (T) gene that is expressed only in postmitotic cells. T-brain-1 (TBR-1) mRNA is largely restricted to the cerebral cortex, where, during embryogenesis, it defines different regions that give rise to the palecortex, limbic cortex and neocortex. TBR-1, Pax-6 and Emx-1 are expressed in the mouse and chicken pallium. The pallio-subpallial boundary lies at the interface between the TBR-1 and Dlx-2 expression domains. Chicken genes homologous to these mouse genes are expressed in topologically comparable patterns during development, suggesting that mouse and chicken may have similar histogenetic specification processes and field homologies. CASK/LIN-2, a membrane-associated guanylate kinase, is required for EGFR localization and signaling. In adult rat brain, CASK is concentrated at neuronal synapses and binds to the cell-surface proteins. CASK can interact with TBR-1, which is involved in forebrain development. CASK enters into the nucleus and binds to a specific DNA sequence (the T-element) in a complex with TBR-1. Thus, CASK acts as a coactivator of TBR-1 to induce transcription of T-element containing genes, including reelin.

Immunogen: Synthetic peptide within Human TBR1 aa 30-75 / 682.

Positive control: Human brain tissue lysates, mouse brain tissue lysates, rat brain tissue lysates, human brain tissue, mouse brain tissue, rat brain tissue, mouse hippocampus tissue, rat hippocampus tissue, E14.5 mouse embryonic brain tissue.

Subcellular location: Nucleus.

Database links: SwissProt: Q16650 Human | Q64336 Mouse

Entrez Gene: 680427 Rat

Recommended Dilutions:

WB 1:1,000

IHC-P 1:500-1:2,000

IF-Tissue 1:500

IHC-Fr 1:500-1:1,000

Storage Buffer: PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images

Fig1: Application: IHC-Fr


Species: Mouse

Site: Cerebral cortex (P0)

Sample: Frozen section

Antibody concentration: 1:1,000

Antigen retrieval: Not required

Fig2: Application: IHC-Fr


Species: Mouse

Site: E14.5 embryonic brain

Sample: Frozen section

Antibody concentration: 1:500

Antigen retrieval: Not required

Fig3: Application: IF-tissue


Species: Mouse

Site: E14.5 embryonic brain

Sample: Paraffin-embedded section

Antibody concentration: 1:500

Fig4: Immunofluorescence analysis of paraffin-embedded mouse brain tissue labeling TBR1 with Rabbit anti-TBR1 antibody (HA750366) at 1/500 dilution.



The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS, and then probed with the primary antibody (HA750366, green) at 1/500 dilution overnight at 4 °C, washed with PBS.

Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Fig5: Western blot analysis of TBR1 on different lysates with Rabbit anti-TBR1 antibody (HA750366) at 1/1,000 dilution.



Lane 1: Human brain tissue lysate (40 µg/Lane)
 Lane 2: Mouse brain tissue lysate (40 µg/Lane)
 Lane 3: Rat brain tissue lysate (40 µg/Lane)

Predicted band size: 74 kDa
 Observed band size: 74 kDa

Exposure time: 1 minute; ECL: K1801;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750366) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.




Fig6: Immunohistochemical analysis of paraffin-embedded mouse embryo tissue with Rabbit anti-TBR1 antibody (HA750366) at 1/2,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750366) at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig7: Immunohistochemical analysis of paraffin-embedded human brain tissue with Rabbit anti-TBR1 antibody (HA750366) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750366) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huabio Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn




Fig8: Immunohistochemical analysis of paraffin-embedded mouse brain tissue with Rabbit anti-TBR1 antibody (HA750366) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750366) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig9: Immunohistochemical analysis of paraffin-embedded rat brain tissue with Rabbit anti-TBR1 antibody (HA750366) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750366) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig10: Immunohistochemical analysis of paraffin-embedded mouse hippocampus tissue with Rabbit anti-TBR1 antibody (HA750366) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750366) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Hangzhou Huan Biotechnology Co.,Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn




Fig11: Immunohistochemical analysis of paraffin-embedded rat hippocampus tissue with Rabbit anti-TBR1 antibody (HA750366) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA750366) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Background References

1. Crespo I et al. Tbr1 Misexpression Alters Neuronal Development in the Cerebral Cortex. *Mol Neurobiol*. 2022 Sep
2. Sollis E et al. Characterization of the TBR1 interactome: variants associated with neurodevelopmental disorders disrupt novel protein interactions. *Hum Mol Genet*. 2023 Apr

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
www.huabio.cn