

# Anti-Parvalbumin Antibody [JM100-08] - BSA and Azide free

## HA750371



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat, Cynomolgus monkey, Pig
<b>Applications:</b>	WB, IHC-P, IHC-Fr, IF-Tissue, IP
<b>Molecular Wt:</b>	Predicted band size: 12 kDa
<b>Clone number:</b>	JM100-08

**Description:** Parvalbumin (PV) is a calcium-binding protein with low molecular weight (typically 9-11 kDa). In humans, it is encoded by the PVALB gene. It is not a member of the albumin family; it is named for its size (parv-, from Latin parvus small) and its ability to coagulate. It has three EF hand motifs and is structurally related to calmodulin and troponin C. Parvalbumin is found in fast-contracting muscles, where its levels are highest, as well as in the brain and some endocrine tissues. Parvalbumin is a small, stable protein containing EF-hand type calcium binding sites. It is involved in calcium signaling. Calcium binding proteins like parvalbumin play a role in many physiological processes, namely cell-cycle regulation, second messenger production, muscle contraction, organization of microtubules and phototransduction. Therefore, calcium-binding proteins must distinguish calcium in the presence of high concentrations of other metal ions. The mechanism for the calcium selectivity has been extensively studied.

**Immunogen:** Recombinant protein within Human Parvalbumin aa 1-110 / 110.

**Positive control:** RPMI 8226 cell lysate, human kidney tissue, mouse kidney tissue, mouse cerebellum tissue, rat cerebellum tissue.

**Subcellular location:** Axon, cytoplasm, synapse.

**Database links:** SwissProt: P20472 Human | P32848 Mouse | P02625 Rat

**Recommended Dilutions:**

<b>WB</b>	1:1,000
<b>IHC-P</b>	1:200-1:1,000
<b>IHC-Fr</b>	1:500-1:1,000
<b>IF-Tissue</b>	1:500
<b>IP</b>	1-2µg/sample

**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4℃ after thawing. Aliquot store at -20℃ or -80℃. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

## Images

**Fig1:** Application: IHC-Fr


Species: Mouse

Site: Cerebellum

Sample: Frozen section

Antibody concentration: 1/500 (Parvalbumin, HA750371, red);  
1/500 (SLC32A1 / VGAT, HA601384, green)

Antigen retrieval: Not required

**Fig2:** Application: IHC-Fr


Species: Rat

Site: Cerebellum

Sample: Frozen section

Antibody concentration: 1/500 (Parvalbumin, HA750371, red);  
1/500 (SLC32A1 / VGAT, HA601384, green)

Antigen retrieval: Not required

**Fig3:** Application: IHC-Fr


Species: Mouse

Site: Cerebellum

Sample: Frozen section

Antibody concentration: 1/1,000

Antigen retrieval: Not required



**Fig4:** Application: IHC-Fr


Species: Rat

Site: Cerebellum

Sample: Frozen section


Antibody concentration: 1/500

Antigen retrieval: Not required




**Fig5:** Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue with Rabbit anti-Parvalbumin antibody (HA750371) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750371) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig6:** Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue with Rabbit anti-Parvalbumin antibody (HA750371) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750371) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig7:** Immunohistochemical analysis of paraffin-embedded human kidney tissue with Rabbit anti-Parvalbumin antibody (HA750371) at 1/200 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750371) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig8:** Immunohistochemical analysis of paraffin-embedded mouse kidney tissue with Rabbit anti-Parvalbumin antibody (HA750371) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750371) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig9:** Western blot analysis of Parvalbumin on different lysates with Rabbit anti-Parvalbumin antibody (HA750371) at 1/1,000 dilution.




Lane 1: RPMI 8226 cell lysate (20 µg/Lane)  
 Lane 2: SK-MEL-28 cell lysate (negative) (20 µg/Lane)  
 Lane 3: Mouse skeletal muscle tissue lysate (20 µg/Lane)  
 Lane 4: Mouse cerebellum tissue lysate (40 µg/Lane)  
 Lane 5: Rat cerebellum tissue lysate (40 µg/Lane)

Predicted band size: 12 kDa  
 Observed band size: 14 kDa  
 Exposure time: Lane 1-3: 30 seconds; Lane 4-5: 3 minutes; ECL: K1801;  
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750371) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.






**Fig10:** Application: IF-tissue

Species: Mouse

Site: Cerebellum

Sample: Paraffin-embedded section

Antibody concentration: 1/500



**Fig11:** Parvalbumin was immunoprecipitated from 0.2 mg RPMI 8226 cell lysate with HA750371 at 2  $\mu$ g/25  $\mu$ l agarose. Western blot was performed from the immunoprecipitate using HA750371 at 1/1,000 dilution. Anti-Rabbit IgG for IP Nano-secondary antibody (NBI01H) at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: RPMI 8226 cell lysate (input)

Lane 2: HA750371 IP in RPMI 8226 cell lysate

Lane 3: Rabbit IgG instead of HA750371 in RPMI 8226 cell lysate

Blocking/Dilution buffer: 5% NFDM/TBST

Exposure time: 10 seconds; ECL: K1801

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

## Background References

1. Cornez G et al. Anatomically discrete sex differences in neuroplasticity in zebra finches as reflected by perineuronal nets. PLoS One 10:e0123199 (2015).
2. Whissell PD et al. Comparative density of CCK- and PV-GABA cells within the cortex and hippocampus. Front Neuroanat 9:124 (2015).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUAABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation