

# Anti-EAAT1 Antibody [JA30-35] - BSA and Azide free

HA750415



**Product Type:** Recombinant Rabbit monoclonal IgG, primary antibodies

**Species reactivity:** Human, Mouse, Rat, Cynomolgus monkey, Pig

**Applications:** WB, IHC-P, IF-Tissue, IHC-Fr

**Molecular Wt:** Predicted band size: 60 kDa

**Clone number:** JA30-35

**Description:** Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate. Functions as a symporter that transports one amino acid molecule together with two or three Na<sup>+</sup> ions and one proton, in parallel with the counter-transport of one K<sup>+</sup> ion. Mediates Cl<sup>-</sup> flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na<sup>+</sup> symport. Plays a redundant role in the rapid removal of released glutamate from the synaptic cleft, which is essential for terminating the postsynaptic action of glutamate. This gene encodes a member of a member of a high affinity glutamate transporter family. This gene functions in the termination of excitatory neurotransmission in central nervous system. Mutations are associated with episodic ataxia, Type 6. Alternative splicing results in multiple transcript variants.

**Immunogen:** Synthetic peptide within Human EAAT1 aa 171-220 / 542.

**Positive control:** Human cerebellum tissue, mouse cerebellum tissue, rat cerebellum tissue, Mouse brain tissue lysate, Mouse cerebellum tissue lysate, Rat brain tissue lysate, Rat cerebellum tissue lysate.

**Subcellular location:** Cell membrane.

**Database links:** SwissProt: P43003 Human | P56564 Mouse | P24942 Rat

**Recommended Dilutions:**

WB 1:5,000

IHC-P 1:1,000-1:8,000

IF-Tissue 1:500

IHC-Fr 1:500

**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

Images

**Fig1: Application: IHC-Fr**



Species: Mouse


Site: Cerebellum

Sample: Frozen section

Antibody concentration: 1:500

Antigen retrieval: Not required

**Fig2: Application: IHC-Fr**



Species: Rat


Site: Cerebellum

Sample: Frozen section

Antibody concentration: 1:500

Antigen retrieval: Not required

**Fig3: Application: IF-tissue**




Species: Mouse

Site: Cerebellum

Sample: Paraffin-embedded section

Antibody concentration: 1:500

**Fig4: Application: IF-tissue**



Species: Rat

Site: Cerebellum

Sample: Paraffin-embedded section

Antibody concentration: 1:500


**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物  
HUABIO  
[www.huabio.cn](http://www.huabio.cn)




**Fig5:** Immunohistochemical analysis of paraffin-embedded human cerebellum tissue with Rabbit anti-EAAT1 antibody (HA750415) at 1/8,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750415) at 1/8,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig6:** Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue with Rabbit anti-EAAT1 antibody (HA750415) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750415) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig7:** Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue with Rabbit anti-EAAT1 antibody (HA750415) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750415) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig8:** Western blot analysis of EAAT1 on different lysates with Rabbit anti-EAAT1 antibody (HA750415) at 1/5,000 dilution.



Lane 1: Mouse brain tissue lysate (no heat)  
 Lane 2: Mouse cerebellum tissue lysate  
 Lane 3: Mouse brain tissue lysate treated with deglycosylation  
 Lane 4: Mouse brain tissue lysate  
 Lane 5: Rat brain tissue lysate (no heat)  
 Lane 6: Rat cerebellum tissue lysate

Notice: no heat means the lysate is not boiled.

Lysates/proteins at 10  $\mu$ g/Lane.

Predicted band size: 60 kDa  
 Observed band size: 60-250 kDa

Exposure time: 25 seconds; ECL: K1801;  
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750415) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Xu NJ et al. Morphine withdrawal increases glutamate uptake and surface expression of glutamate transporter GLT1 at hippocampal synapses. *J Neurosci*. 23(11): 4775-4784 (2003).
2. Ueda, Hideho et al. Caveolin-1 Localization in Müller Cells of the Retina *ACTA HISTOCHEMICA ET CYTOCHEMICA*. 35: 423-428 (2002).

**Hangzhou Huan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
www.huabio.cn