

Anti-CXCR3 Antibody [JA61-33] - BSA and Azide free

HA750422



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	WB
Molecular Wt:	Predicted band size: 41 kDa
Clone number:	JA61-33

Description: Chemokine receptor CXCR3 is a Gai protein-coupled receptor in the CXC chemokine receptor family. Other names for CXCR3 are G protein-coupled receptor 9 (GPR9) and CD183. CXCR3 is able to regulate leukocyte trafficking. Binding of chemokines to CXCR3 induces various cellular responses, most notably integrin activation, cytoskeletal changes and chemotactic migration. CXCR3-ligand interaction attracts Th1 cells and promotes Th1 cell maturation. As a consequence of chemokine-induced cellular desensitization (phosphorylation-dependent receptor internalization), cellular responses are typically rapid and short in duration. Cellular responsiveness is restored after dephosphorylation of intracellular receptors and subsequent recycling to the cell surface. A hallmark of CXCR3 is its prominent expression in in vitro cultured effector/memory T cells, and in T cells present in many types of inflamed tissues. In addition, CXCL9, CXCL10 and CXCL11 are commonly produced by local cells in inflammatory lesion, suggesting that CXCR3 and its chemokines participate in the recruitment of inflammatory cells. Additionally, CXCR3 has been implicated in wound healing.

Immunogen:	Recombinant protein within human CXCR3 aa 278-298 (Extracellular)/368.
Positive control:	HeLa cell lysate, HepG2 cell lysate, PANC-1 cell lysate, K-562 cell lysate, HT-29 cell lysate, Human lung tissue lysate.
Subcellular location:	Cell membrane.
Database links:	SwissProt: P49682 Human
Recommended Dilutions:	
WB	1:1,000-1:5,000
Storage Buffer:	1*PBS (pH7.4).
Storage Instruction:	Store at +4℃ after thawing. Aliquot store at -20℃ or -80℃. Avoid repeated freeze / thaw cycles.
Purity:	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of CXCR3 on different lysates with Rabbit anti-CXCR3 antibody (HA750422) at 1/5,000 dilution.

Lane 1: HeLa cell lysate
Lane 2: HepG2 cell lysate
Lane 3: PANC-1 cell lysate
Lane 4: K-562 cell lysate
Lane 5: HT-29 cell lysate



Lysates/proteins at 20 µg/Lane.

Predicted band size: 41 kDa
Observed band size: 50 kDa


Exposure time: 1 minute; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750422) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of CXCR3 on different lysates with Rabbit anti-CXCR3 antibody (HA750422) at 1/1,000 dilution.

Lane 1: Human lung tissue lysate (20 µg/Lane)
Lane 2: K-562 cell lysate (10 µg/Lane)



Predicted band size: 41 kDa
Observed band size: 50 kDa

Exposure time: 57 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750422) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Tanaka A et al. Th2 and regulatory immune reactions contribute to IgG4 production and the initiation of Mikulicz disease. *Arthritis Rheum* 64:254-63 (2012).
2. Song CH et al. Identification of chemoattractive factors involved in the migration of bone marrow-derived mesenchymal stem cells to brain lesions caused by prions. *J Virol* 85:11069-78 (2011).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn