

# Anti-STING Antibody [JM03-47] - BSA and Azide free

## HA750437



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Rat
<b>Applications:</b>	WB, IHC-P, IF-Cell, FC, IF-Tissue, IP
<b>Molecular Wt:</b>	Predicted band size: 42 kDa
<b>Clone number:</b>	JM03-47

<b>Description:</b>	Facilitator of innate immune signaling that promotes the production of type I interferon (IFN-alpha and IFN-beta). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm. Able to activate both NF-kappa-B and IRF3 transcription pathways to induce expression of type I interferon and exert a potent anti-viral state following expression. May be involved in translocon function, the translocon possibly being able to influence the induction of type I interferons. May be involved in transduction of apoptotic signals via its association with the major histocompatibility complex class II (MHC-II). Mediates death signaling via activation of the extracellular signal-regulated kinase (ERK) pathway.
<b>Immunogen:</b>	Recombinant protein within human STING aa 117-379.
<b>Positive control:</b>	THP-1 cell lysate, SW620 cell lysate, human lung tissue lysate, THP-1, human tonsil tissue, rat lung tissue.
<b>Subcellular location:</b>	Endoplasmic reticulum membrane, Cytoplasm, perinuclear region, Endoplasmic reticulum-Golgi intermediate compartment membrane, Golgi apparatus membrane, Cytoplasmic vesicle, autophagosome membrane, Mitochondrion outer membrane, Cell membrane.
<b>Database links:</b>	SwissProt: Q86WW6 Human   F1M391 Rat
<b>Recommended Dilutions:</b>	
WB	1:1,000
IHC-P	1:1,000
IF-Cell	1:100
FC	1:1,000
IF-Tissue	1:200
IP	1-2µg/sample
<b>Storage Buffer:</b>	PBS (pH7.4).
<b>Storage Instruction:</b>	Store at +4℃ after thawing. Aliquot store at -20℃ or -80℃. Avoid repeated freeze / thaw cycles.
<b>Purity:</b>	Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation


## Images

**Fig1:** Western blot analysis of STING on different lysates with Rabbit anti-STING antibody (HA750437) at 1/1,000 dilution.

Lane 1: THP-1 cell lysate (15 µg/Lane)

Lane 2: SW620 cell lysate (15 µg/Lane)

Lane 3: Human lung tissue lysate (30 µg/Lane)



Predicted band size: 42 kDa


Observed band size: 35 kDa

Exposure time: 1 minute 2 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750437) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Immunocytochemistry analysis of THP-1 cells labeling STING with Rabbit anti-STING antibody (HA750437) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-STING antibody (HA750437) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation




**Fig3:** Immunohistochemical analysis of paraffin-embedded human tonsil tissue with Rabbit anti-STING antibody (HA750437) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750437) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig4:** Immunohistochemical analysis of paraffin-embedded rat lung tissue with Rabbit anti-STING antibody (HA750437) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750437) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig5:** Flow cytometric analysis of THP-1 cells labeling STING.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA750437, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).



**Fig6:** STING was immunoprecipitated from 0.2 mg THP-1 cell lysate with HA750437 at 2  $\mu$ g/10  $\mu$ l beads. Western blot was performed from the immunoprecipitate using HA750437 at 1/1,000 dilution. Anti-Rabbit IgG for IP Nano-secondary antibody (NBI01H) at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: THP-1 cell lysate (input)

Lane 2: HA750437 IP in THP-1 cell lysate

Lane 3: Rabbit IgG instead of HA750437 in THP-1 cell lysate

Blocking/Dilution buffer: 5% NFDM/TBST

Exposure time: 3 minutes; ECL: K1801

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Panchanathan R et al. Identification of a negative feedback loop between cyclic di-GMP-induced levels of IFI16 and p202 cytosolic DNA sensors and STING. *Innate Immun* 20(7):751-9 (2014).
2. Orzalli MH et al. Nuclear interferon-inducible protein 16 promotes silencing of herpesviral and transfected DNA. *Proc Natl Acad Sci U S A* 110:E4492-501 (2013).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUAABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation