

# Anti-F4/80 Antibody [PSH0-85] - BSA and Azide free

## HA750665



**Product Type:** Recombinant Rabbit monoclonal IgG, primary antibodies

**Species reactivity:** Mouse, Rat

**Applications:** WB, IHC-P, IF-Tissue, IHC-Fr

**Molecular Wt:** Predicted band size: 102 kDa

**Clone number:** PSH0-85

**Description:** EGF-like module-containing mucin-like hormone receptor-like 1 also known as F4/80 is a protein encoded by the ADGRE1 gene. EMR1 is a member of the adhesion GPCR family. Adhesion GPCRs are characterized by an extended extracellular region often possessing N-terminal protein modules that is linked to a TM7 region via a domain known as the GPCR-Autoproteolysis INducing (GAIN) domain. EMR1 expression in human is restricted to eosinophils and is a specific marker for these cells. The murine homolog of EMR1, F4/80, is a well-known and widely used marker of murine macrophage populations. The N-terminal fragment (NTF) of EMR1 contains 4-6 Epidermal Growth Factor-like (EGF-like) domains in human and 4-7 EGF-like domains in the mouse. Utilizing F4/80 knockout mice, Lin et al. showed that F4/80 is not necessary for the development of tissue macrophages but is required for the induction of efferent CD8+ regulatory T cells needed for peripheral tolerance.

**Immunogen:** Recombinant protein within mouse F4/80 aa 1-650 / 931.

**Positive control:** Mouse kidney tissue lysate, RAW264.7 cell lysate, Rat kidney tissue lysate, Rat spleen tissue lysate, mouse liver tissue, mouse spleen tissue, rat liver tissue, rat spleen tissue, mouse lung tissue, rat lung tissue.

**Subcellular location:** Cell membrane.

**Database links:** SwissProt: Q61549 Mouse | Q5Y4N8 Rat

**Recommended Dilutions:**

**WB** 1:1,000

**IHC-P** 1:200-1:1,000

**IF-Tissue** 1:500

**IHC-Fr** 1:1,000

**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of F4/80 on different lysates with Rabbit anti-F4/80 antibody (HA750665) at 1/1,000 dilution.



Lane 1: Mouse kidney tissue lysate (no heat) (40 µg/Lane)  
 Lane 2: Mouse kidney tissue lysate (40 µg/Lane)  
 Lane 3: RAW264.7 cell lysate (no heat) (20 µg/Lane)  
 Lane 4: Rat kidney tissue lysate (no heat) (40 µg/Lane)

Notice: no heat means the lysate is not boiled.

Predicted band size: 102 kDa  
 Observed band size: 160 kDa

Exposure time: 59 seconds; ECL: K1801;  
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750665) at 1/1,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.


**Fig2:** Western blot analysis of F4/80 on different lysates with Rabbit anti-F4/80 antibody (HA750665) at 1/1,000 dilution.

Lane 1: RAW264.7 cell lysate (no heat) (20 µg/Lane)  
 Lane 2: L929 cell lysate (no heat) (negative) (20 µg/Lane)  
 Lane 3: Rat spleen tissue lysate (70°C heat) (40 µg/Lane)

Notice: no heat means the lysate is not boiled.

Predicted band size: 102 kDa  
 Observed band size: 160 kDa

Exposure time: 2 minutes;  
 4-20% SDS-PAGE gel.



Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750665) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1:100,000 dilution was used for 1 hour at room temperature.


**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

**Fig3: Application: IHC-Fr**


Species: Mouse

Site: Spleen

Sample: Frozen section

Antibody concentration: 1/1,000 (F4/80, HA750665, rabbit, red);  
1/500 (Iba1, HA601376, Chicken, green)

Antigen retrieval: Not required

**Fig4: Application: IHC-Fr**


Species: Rat

Site: Spleen

Sample: Frozen section

Antibody concentration: 1/1,000 (F4/80, HA750665, rabbit, red);  
1/500 (Iba1, HA601376, Chicken, green)

Antigen retrieval: Not required

**Fig5: Immunohistochemical analysis of paraffin-embedded mouse liver tissue with Rabbit anti-F4/80 antibody (HA750665) at 1/1,000 dilution.**

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750665) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


**Hangzhou Huan Biotechnology Co.,Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn




**Fig6:** Immunohistochemical analysis of paraffin-embedded mouse spleen tissue with Rabbit anti-F4/80 antibody (HA750665) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750665) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig7:** Immunohistochemical analysis of paraffin-embedded rat liver tissue with Rabbit anti-F4/80 antibody (HA750665) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750665) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig8:** Immunohistochemical analysis of paraffin-embedded rat spleen tissue with Rabbit anti-F4/80 antibody (HA750665) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750665) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




**Fig9:** Immunohistochemical analysis of paraffin-embedded mouse lung tissue with Rabbit anti-F4/80 antibody (HA750665) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750665) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig10:** Immunohistochemical analysis of paraffin-embedded rat lung tissue with Rabbit anti-F4/80 antibody (HA750665) at 1/2,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750665) at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig11:** Application: IF-tissue

Species: Mouse

Site: Liver

Sample: Paraffin-embedded section

Antibody concentration: 1/500


**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications: WB=Western blot IHC=P=Immunohistochemistry (paraffin) IF=Cell=Immunofluorescence (Cell) IF=Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation


**Fig12:** Application: IF-tissue

Species: Rat

Site: Liver

Sample: Paraffin-embedded section

Antibody concentration: 1/500


**Fig13:** Application: IF-tissue

Species: Mouse

Site: spleen

Sample: Paraffin-embedded section

Antibody concentration: 1/500

**Fig14:** Application: IF-tissue

Species: Mouse

Site: small intestine

Sample: Paraffin-embedded section

Antibody concentration: 1/500

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Deng R et al. Periosteal CD68+ F4/80+ Macrophages Are Mechanosensitive for Cortical Bone Formation by Secretion and Activation of TGF- $\beta$ 1. *Adv Sci (Weinh)*. 2022 Jan
2. Shin AE et al. F4/80+Ly6Chigh Macrophages Lead to Cell Plasticity and Cancer Initiation in Colitis. *Gastroenterology*. 2023 Apr

**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn