

Anti-PACT (PKR activating protein) / PRKRA Antibody [PSH02-15] - BSA and Azide free

**HA750785**



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IHC-P, IF-Cell, IF-Tissue, FC
<b>Molecular Wt:</b>	Predicted band size: 34 kDa
<b>Clone number:</b>	PSH02-15

**Description:** PRKRA can activate EIF2AK2/PKR in the absence of double-stranded RNA (dsRNA), leading to phosphorylation of EIF2S1/EF12-alpha and inhibition of translation and induction of apoptosis. Required for siRNA production by DICER1 and for subsequent siRNA-mediated post-transcriptional gene silencing. Does not seem to be required for processing of pre-miRNA to miRNA by DICER1. Promotes UBC9-p53/TP53 association and sumoylation and phosphorylation of p53/TP53 at 'Lys-386' at 'Ser-392' respectively and enhances its activity in a EIF2AK2/PKR-dependent manner.

**Immunogen:** Recombinant protein within human PACT aa 1-313 (O75569).

**Positive control:** 293T cell lysate, K-562 cell lysate, Jurkat cell lysate, HepG2 cell lysate, MCF7 cell lysate, Hela cell lysate, HL-60 cell lysate, C2C12 cell lysate, PC-12 cell lysate, mouse testis tissue lysate, mouse liver tissue lysate, rat testis tissue lysate, rat brain tissue lysate, human testis tissue, rat brain tissue, rat cerebral cortex tissue, rat hippocampus tissue, 293T, PC-12.

**Subcellular location:** Cytoplasm, perinuclear region

**Database links:** SwissProt: O75569 Human | Q9WTX2 Mouse | Q4V8C7 Rat

**Recommended Dilutions:**

<b>WB</b>	1:1,000
<b>IHC-P</b>	1:1,000-1:5,000
<b>IF-Cell</b>	1:100
<b>IF-Tissue</b>	1:200
<b>FC</b>	1 µg/mL

**Storage Buffer:** 1\*PBS (pH7.4).

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of PACT (PKR activating protein) / PRKRA on different lysates with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/1,000 dilution.



Lane 1: 293T cell lysate  
 Lane 2: K-562 cell lysate  
 Lane 3: Jurkat cell lysate  
 Lane 4: HepG2 cell lysate  
 Lane 5: MCF7 cell lysate  
 Lane 6: Hela cell lysate  
 Lane 7: HL-60 cell lysate  
 Lane 8: C2C12 cell lysate  
 Lane 9: PC-12 cell lysate  
 Lane 10: Mouse testis tissue lysate  
 Lane 11: Mouse liver tissue lysate  
 Lane 12: Rat testis tissue lysate  
 Lane 13: Rat brain tissue lysate


Cell lysates: 20ug/lane, tissue lysates: 40ug/lane

Predicted band size: 34 kDa  
 Observed band size: 34 kDa

Exposure time: 1 minutes 2 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750785) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunohistochemical analysis of paraffin-embedded human testis tissue with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750785) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

华安生物  
 HUABIO  
 www.huabio.cn




**Fig3:** Immunohistochemical analysis of paraffin-embedded rat brain tissue with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/5,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750785) at 1/5,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig4:** Immunofluorescence analysis of paraffin-embedded rat cerebral cortex tissue labeling PACT (PKR activating protein) / PRKRA with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/200 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS, and then probed with the primary antibody (HA750785, green) at 1/200 dilution overnight at 4 °C, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).



**Fig5:** Immunofluorescence analysis of paraffin-embedded rat hippocampus tissue labeling PACT (PKR activating protein) / PRKRA with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS, and then probed with the primary antibody (HA750785, green) at 1/200 dilution overnight at 4 °C, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).


**Fig6:** Immunocytochemistry analysis of 293T cells labeling PACT (PKR activating protein) / PRKRA with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.


Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

**Fig7:** Immunocytochemistry analysis of PC-12 cells labeling PACT (PKR activating protein) / PRKRA with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-PACT (PKR activating protein) / PRKRA antibody (HA750785) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



**Fig8:** Flow cytometric analysis of 293T cells labeling PACT (PKR activating protein) / PRKRA.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA750785, 1 $\mu$ g/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4 $^{\circ}$ C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4 $^{\circ}$ C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Pullagura SRN et al. Functional Redundancy of DICER Cofactors TARBP2 and PRKRA During Murine Embryogenesis Does Not Involve miRNA Biogenesis. *Genetics* 208:1513-1522 (2018).
2. Vaughn LS et al. DYT-PRKRA Mutation P222L Enhances PACT's Stimulatory Activity on Type I Interferon Induction. *Biomolecules* 12:N/A (2022).

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn