

# Anti-beta Amyloid 1-40 Antibody [PSH02-82] - BSA and Azide free

HA750789



**Product Type:** Recombinant Rabbit monoclonal IgG, primary antibodies

**Species reactivity:** Human, Mouse, Cynomolgus monkey, Pig

**Applications:** IHC-P, WB

**Molecular Wt:** Predicted band size: 4 kDa

**Clone number:** PSH02-82

**Description:** This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides. Some of these peptides are secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote transcriptional activation, while others form the protein basis of the amyloid plaques found in the brains of patients with Alzheimer disease. In addition, two of the peptides are antimicrobial peptides, having been shown to have bacteriocidal and antifungal activities. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been found for this gene.

**Immunogen:** Synthetic peptide within Human beta Amyloid 1-40 peptide.

**Positive control:** APP/PS1, 6-month mouse of AD brain tissue.

**Subcellular location:** Cell membrane, Membrane, Perikaryon, Cell projection, growth cone, Membrane, clathrin-coated pit, Early endosome, Cytoplasmic vesicle.

**Database links:** SwissProt: P05067 Human | P12023 Mouse

**Recommended Dilutions:**

IHC-P 1:1,000

WB 1:1,000

**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

## Images



**Fig1:** Western blot analysis of beta Amyloid 1-40 on different peptides with Rabbit anti-beta Amyloid 1-40 antibody (HA750789) at 1/1,000 dilution.

Lane 1: Human A $\beta$ 37 full length peptide (negative)  
 Lane 2: Human A $\beta$ 38 full length peptide (negative)  
 Lane 3: Human A $\beta$ 39 full length peptide (negative)  
 Lane 4: Human A $\beta$ 40 full length peptide (positive)  
 Lane 5: Human A $\beta$ 41 full length peptide (negative)  
 Lane 6: Human A $\beta$ 42 full length peptide (negative)  
 Lane 7: Human A $\beta$ 43 full length peptide (negative)  
 Lane 8: Mouse A $\beta$ 40 full length peptide (positive)


Lysates/proteins at 100 ng/Lane.

Predicted band size: 4 kDa  
 Observed band size: 4 kDa

Exposure time: 10 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750789) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunohistochemical analysis of paraffin-embedded APP/PS1, 6-month mouse of AD brain tissue with Rabbit anti-beta Amyloid 1-40 antibody (HA750789) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750789) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn



**Fig3:** Immunohistochemical analysis of paraffin-embedded mouse brain tissue (negative control) with Rabbit anti-beta Amyloid 1-40 antibody (HA750789) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750789) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

#### Background References

1. Janelidze S et al. Head-to-Head Comparison of 8 Plasma Amyloid- $\beta$  42/40 Assays in Alzheimer Disease. *JAMA Neurol.* 2021 Nov

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn