

## Anti-alpha Tubulin Antibody [PSH02-95] - BSA and Azide free

# HA750850



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IHC-P, IF-Cell, FC, IP
<b>Molecular Wt:</b>	Predicted band size: 50 kDa
<b>Clone number:</b>	PSH02-95

**Description:** The cytoskeleton consists of three types of cytosolic fibers: microtubules, microfilaments (actin filaments), and intermediate filaments. Globular tubulin subunits comprise the microtubule building block, with  $\alpha/\beta$ -tubulin heterodimers forming the tubulin subunit common to all eukaryotic cells. Acetylation of alpha chains at Lys-40 stabilizes microtubules and affects affinity and processivity of microtubule motors. This modification has a role in multiple cellular functions, ranging from cell motility, cell cycle progression or cell differentiation to intracellular trafficking and signaling.

**Immunogen:** Synthetic peptide within Human alpha Tubulin aa 402-451 / 451.

**Positive control:** HeLa cell lysate, Daudi cell lysate, Jurkat cell lysate, A431 cell lysate, K-562 cell lysate, 293T cell lysate, NIH/3T3 cell lysate, PC-12 cell lysate, Rat brain tissue lysate, Mouse brain tissue lysate, Mouse spleen tissue lysate, human tonsil tissue, rat brain tissue, Daudi, K-562, HeLa.

**Subcellular location:** Cytoplasm, Cytoskeleton, Microtubule.

**Database links:** SwissProt: P68363 Human | P68368 Mouse | Q5XIF6 Rat

### Recommended Dilutions:

<b>WB</b>	1:1,000-5,000
<b>IHC-P</b>	1:1,000
<b>IF-Cell</b>	1:100
<b>FC</b>	1:1,000
<b>IP</b>	1-2 $\mu$ g/sample

**Storage Buffer:** 1\*PBS (pH7.4).

**Storage Instruction:** Store at +4 $^{\circ}$ C after thawing. Aliquot store at -20 $^{\circ}$ C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1:** Western blot analysis of alpha Tubulin on different lysates with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/5,000 dilution.




Lane 1: HeLa cell lysate (20 µg/Lane)  
 Lane 2: Daudi cell lysate (20 µg/Lane)  
 Lane 3: Jurkat cell lysate (20 µg/Lane)  
 Lane 4: A431 cell lysate (20 µg/Lane)  
 Lane 5: K-562 cell lysate (20 µg/Lane)  
 Lane 6: 293T cell lysate (20 µg/Lane)  
 Lane 7: NIH/3T3 cell lysate (20 µg/Lane)  
 Lane 8: PC-12 cell lysate (20 µg/Lane)  
 Lane 9: Rat brain tissue lysate (40 µg/Lane)  
 Lane 10: Mouse brain tissue lysate (40 µg/Lane)  
 Lane 11: Mouse spleen tissue lysate (40 µg/Lane)

Predicted band size: 50 kDa  
 Observed band size: 50 kDa

Exposure time: 3 minutes; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA750850) at 1/5,000 dilution was used in 5% NFDm/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.



**Fig2:** Immunofluorescence analysis of paraffin-embedded human tonsil tissue labeling alpha Tubulin with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS, and then probed with the primary antibody (HA750850, green) at 1/1,000 dilution overnight at 4 °C, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物  
 HUABIO  
 www.huabio.cn



**Fig3:** Immunohistochemical analysis of paraffin-embedded rat brain tissue with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) (high pressure) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750850) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


**Fig4:** Immunocytochemistry analysis of Daudi cells labeling alpha Tubulin with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.


Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

**Fig5:** Immunocytochemistry analysis of K-562 cells labeling alpha Tubulin with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/100 dilution.




Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-alpha Tubulin antibody (HA750850) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



**Fig6:** Flow cytometric analysis of HeLa cells labeling alpha Tubulin.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA750850, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).



**Fig7:** alpha Tubulin was immunoprecipitated from 0.2 mg HeLa cell lysate with HA750850 at 2 µg/10 µl beads. Western blot was performed from the immunoprecipitate using HA750850 at 1/1,000 dilution. Anti-Rabbit IgG for IP Nano-secondary antibody (NBI01H) at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: HeLa cell lysate (input)  
Lane 2: HA750850 IP in HeLa cell lysate  
Lane 3: Rabbit IgG instead of HA750850 in HeLa cell lysate

Blocking/Dilution buffer: 5% NFDm/TBST  
Exposure time: 2 seconds; ECL: K1801

---

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

---

### Background References

1. Girão H et al. alpha-tubulin detyrosination fine-tunes kinetochore-microtubule attachments. *Nat Commun.* 2024 Nov
2. Wethekam LC et al. alpha-tubulin regulation by 5' introns in *Saccharomyces cerevisiae*. *Genetics.* 2023 Dec

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物  
HUA B I O  
www.huabio.cn