

# Anti-TPH2 Antibody [PSH04-89] - BSA and Azide free

HA750976



**Product Type:** Recombinant Rabbit monoclonal IgG, primary antibodies

**Species reactivity:** Mouse, Rat, Cynomolgus monkey, Pig

**Applications:** WB, IHC-P, IHC-Fr, IP

**Molecular Wt:** Predicted band size: 56 kDa

**Clone number:** PSH04-89

**Description:** Tryptophan hydroxylase 2 (TPH2) is an isozyme of tryptophan hydroxylase found in vertebrates. In humans, TPH2 is primarily expressed in the serotonergic neurons of the brain, with the highest expression in the raphe nucleus of the midbrain. Until the discovery of TPH2 in 2003, serotonin levels in the central nervous system were believed to be regulated by serotonin synthesis in peripheral tissues, in which tryptophan hydroxylase is the dominant form.

**Immunogen:** Recombinant protein with Human TPH2 aa 1-490.

**Positive control:** Mouse hippocampus tissue lysate, mouse brain tissue lysate, rat brain tissue lysate, rat hippocampus tissue lysate, mouse midbrain tissue, mouse cerebrum tissue, rat cerebrum tissue, mouse brain (raphe nucleus) tissue.

**Subcellular location:** Cytosol, neuron projection.

**Database links:** SwissProt: Q8CGV2 Mouse | Q8CGU9 Rat

**Recommended Dilutions:**

WB 1:2,000

IHC-P 1:500

IHC-Fr 1:500

IP 1-2 $\mu$ g/sample

**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.


Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

## Images

**Fig1: Application: IHC-Fr**

Species: Mouse

Site: Raphe nucleus


Sample: Frozen section

Antibody concentration: 1/500

Antigen retrieval: Not required

**Fig2: Immunohistochemical analysis of paraffin-embedded mouse midbrain tissue with Rabbit anti-TPH2 antibody (HA750976) at 1/500 dilution.**

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750976) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

**Fig3: Immunohistochemical analysis of paraffin-embedded mouse cerebrum tissue with Rabbit anti-TPH2 antibody (HA750976) at 1/500 dilution.**

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750976) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物  
HUABIO  
[www.huabio.cn](http://www.huabio.cn)



**Fig4:** Immunohistochemical analysis of paraffin-embedded rat cerebrum tissue with Rabbit anti-TPH2 antibody (HA750976) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH<sub>2</sub>O and PBS, and then probed with the primary antibody (HA750976) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



**Fig5:** Western blot analysis of TPH2 on different lysates with Rabbit anti-TPH2 antibody (HA750976) at 1/2,000 dilution.

Lane 1: Mouse brain tissue lysate  
 Lane 2: Mouse kidney tissue lysate (negative)  
 Lane 3: Rat brain tissue lysate  
 Lane 4: Rat kidney tissue lysate (negative)

Lysates/proteins at 20 µg/Lane.

Predicted band size: 56 kDa  
 Observed band size: 50 kDa

Exposure time: 1 minute 2 seconds; ECL: K1801;  
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA750976) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.


Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn



**Fig6:** TPH2 was immunoprecipitated from 0.2 mg rat brain tissue lysate with HA750976 at 2  $\mu$ g/10  $\mu$ l beads. Western blot was performed from the immunoprecipitate using HA750976 at 1/1,000 dilution. Anti-Rabbit IgG for IP Nano-secondary antibody (NBI01H) at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: rat brain tissue lysate (input)  
 Lane 2: HA750976 IP in rat brain tissue lysate  
 Lane 3: Rabbit IgG instead of HA750976 in rat brain tissue lysate

Blocking/Dilution buffer: 5% NFDM/TBST

Exposure time: 3 minutes; ECL: K1801

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

### Background References

1. Zaniewska M et al. Tph2 Gene Expression Defines Ethanol Drinking Behavior in Mice. *Cells*. 2022 Mar
2. Liu H et al. TPH2 in the Dorsal Raphe Nuclei Regulates Energy Balance in a Sex-Dependent Manner. *Endocrinology*. 2021 Jan

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn