

Anti-clAP1 Antibody [PSH06-71] - BSA and Azide free HA751100



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat
Applications:	WB
Molecular Wt:	Predicted band size: 70 kDa
Clone number:	PSH06-71

Description: Baculoviral IAP repeat-containing protein 2 (also known as clAP1) is a protein that in humans is encoded by the BIRC2 gene. clAP1 is a member of the Inhibitor of Apoptosis family that inhibit apoptosis by interfering with the activation of caspases.

Immunogen: Recombinant protein within human clAP1 aa 201-618.

Positive control: HT-29 cell lysate, TF-1 cell lysate, HepG2 cell lysate, HeLa cell lysate, THP-1 cell lysate, Jurkat cell lysate, A549 cell lysate, HEK-293 cell lysate, NIH/3T3 cell lysate, PC-12 cell lysate, Mouse testis tissue lysate, Mouse brain tissue lysate, Rat testis tissue lysate, Rat brain tissue lysate.

Subcellular location: Cytoplasm, Nucleus.

Database links: SwissProt: Q13490 Human | Q62210 Mouse
Entrez Gene: 60371 Rat

Recommended Dilutions:
WB 1:2,000

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of cIAP1 on different lysates with Rabbit anti-cIAP1 antibody (HA751100) at 1/2,000 dilution.



Lane 1: HT-29 cell lysate (20 µg/Lane)
 Lane 2: TF-1 cell lysate (20 µg/Lane)
 Lane 3: HepG2 cell lysate (20 µg/Lane)
 Lane 4: HeLa cell lysate (20 µg/Lane)
 Lane 5: THP-1 cell lysate (20 µg/Lane)
 Lane 6: Jurkat cell lysate (20 µg/Lane)
 Lane 7: A549 cell lysate (20 µg/Lane)
 Lane 8: HEK-293 cell lysate (20 µg/Lane)
 Lane 9: NIH/3T3 cell lysate (20 µg/Lane)
 Lane 10: PC-12 cell lysate (20 µg/Lane)
 Lane 11: Mouse testis tissue lysate (40 µg/Lane)
 Lane 12: Mouse brain tissue lysate (40 µg/Lane)
 Lane 13: Rat testis tissue lysate (40 µg/Lane)
 Lane 14: Rat brain tissue lysate (40 µg/Lane)


Predicted band size: 70 kDa

Observed band size: 70 kDa

Exposure time: 2 minutes; ECL: K1801;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (HA751100) at 1/2,000 dilution was used in 5% NFDN/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Western blot analysis of cIAP1 on different lysates with Rabbit anti-cIAP1 antibody (HA751100) at 1/2,000 dilution.



Lane 1: HAP1-parental cell lysate (10 µg/Lane)
 Lane 2: HAP1-cIAP1 KD cell lysate (10 µg/Lane)

Predicted band size: 70 kDa

Observed band size: 70 kDa

Exposure time: 78 seconds; ECL: K1801;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (HA751100) at 1/2,000 dilution was used in K1803 at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Zadoroznyj A et al. Cytoplasmic and Nuclear Functions of cIAP1. *Biomolecules*. 2022 Feb
2. Yang X et al. TRIM56 promotes malignant progression of glioblastoma by stabilizing cIAP1 protein. *J Exp Clin Cancer Res*. 2022 Dec

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn