

# Anti-DDIT3 Antibody [PSH07-65] - BSA and Azide free

## HA751178



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB, IF-Cell, FC
<b>Molecular Wt:</b>	Predicted band size: 19 kDa
<b>Clone number:</b>	PSH07-65

**Description:** DNA damage-inducible transcript 3, also known as C/EBP homologous protein (CHOP), is a pro-apoptotic transcription factor that is encoded by the DDIT3 gene. It is a member of the CCAAT/enhancer-binding protein (C/EBP) family of DNA-binding transcription factors. The protein functions as a dominant-negative inhibitor by forming heterodimers with other C/EBP members, preventing their DNA binding activity. The protein is implicated in adipogenesis and erythropoiesis and has an important role in the cell's stress response.

**Immunogen:** Recombinant protein within human DDIT3 aa 1-169.

**Positive control:** HeLa treated with 2µg/mL tunicamycin for 8 hours cell lysate, C2C12 treated with 2µg/mL thapsigargin for 8 hours cell lysate, C6 cell lysate, C6 treated with 2µg/mL tunicamycin for 8 hours cell lysate, HeLa cells treated with 2µg/mL tunicamycin for 8 hours, C2C12 cells treated with 2µg/mL thapsigargin for 8 hours, C6 cells treated with 2µg/mL tunicamycin for 8 hours.

**Subcellular location:** Cytoplasm, Nucleus.

**Database links:** SwissProt: P35638 Human | P35639 Mouse | Q62857 Rat

**Recommended Dilutions:**

<b>WB</b>	1:5,000
<b>IF-Cell</b>	1:100
<b>FC</b>	1:1,000

**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345


Service mail:support@huabio.cn

 华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

## Images

**Fig1:** Western blot analysis of DDIT3 on different lysates with Rabbit anti-DDIT3 antibody (HA751178) at 1/5,000 dilution.



Lane 1: HeLa cell lysate

Lane 2: HeLa treated with 2 $\mu$ g/mL tunicamycin for 8 hours cell lysate

Lane 3: C2C12 cell lysate

Lane 4: C2C12 treated with 2 $\mu$ g/mL thapsigargin for 8 hours cell lysate

Lane 5: C6 cell lysate

Lane 6: C6 treated with 2 $\mu$ g/mL tunicamycin for 8 hours cell lysate

Lysates/proteins at 20  $\mu$ g/Lane.

Predicted band size: 19 kDa


Observed band size: 27 kDa

Exposure time: Lane 1-2: 3 minutes; Lane 3-6: 10 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA751178) at 1/5,000 dilution was used in 5% NFDM/TBST at 4 $^{\circ}$ C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

**Fig2:** Immunocytochemistry analysis of HeLa cells treated with 2 $\mu$ g/mL tunicamycin for 8 hours labeling DDIT3 with Rabbit anti-DDIT3 antibody (HA751178) at 1/100 dilution.



Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-DDIT3 antibody (HA751178) at 1/100 dilution in 1% BSA in PBST overnight at 4 $^{\circ}$ C. Goat Anti-Rabbit IgG H&L (iFluor<sup>TM</sup> 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4 $^{\circ}$ C. Goat Anti-Mouse IgG H&L (iFluor<sup>TM</sup> 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.


Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn


华安生物  
HUABIO  
www.huabio.cn



**Fig3:** Immunocytochemistry analysis of C2C12 cells treated with 2 $\mu$ g/mL thapsigargin for 8 hours labeling DDIT3 with Rabbit anti-DDIT3 antibody (HA751178) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-DDIT3 antibody (HA751178) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.


Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



**Fig4:** Immunocytochemistry analysis of C6 cells treated with 2 $\mu$ g/mL tunicamycin for 8 hours labeling DDIT3 with Rabbit anti-DDIT3 antibody (HA751178) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-DDIT3 antibody (HA751178) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (M1305-2, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.



**Fig5:** Flow cytometric analysis of C2C12 cells treated with 2 $\mu$ g/mL thapsigargin for 8 hours labeling DDIT3.

Cells were fixed and permeabilized. Then stained with the primary antibody (HA751178, 1/1,000) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4°C for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody (HA1121) at 1/1,000 dilution for 30 minutes at +4°C. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).

---

**Note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

---

### Background References

1. Li M et al. DDIT3 Directs a Dual Mechanism to Balance Glycolysis and Oxidative Phosphorylation during Glutamine Deprivation. *Adv Sci (Weinh)*. 2021 Jun
2. Wang Y et al. DDIT3 aggravates pulpitis by modulating M1 polarization through EGR1 in macrophages. *Int Immunopharmacol*. 2023 Jul

**Hangzhou Huaan Biotechnology Co., Ltd.**

Orders: 0086-571-88062880

Technical: 0086-571-89986345

Service mail: support@huabio.cn

 华安生物  
HUAABIO  
www.huabio.cn

Applications: WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation