

Anti-Synapsin I + II Antibody [PSH10-29] - BSA and Azide free

HA751346



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat, Cynomolgus monkey, Pig
Applications:	WB, IF-Cell, IHC-Fr, IHC-P, IP, IF-Tissue
Molecular Wt:	Predicted band size: 74 kDa
Clone number:	PSH10-29

Description: Synapsin I, is the collective name for Synapsin Ia and Synapsin Ib, two nearly identical phosphoproteins that in humans are encoded by the SYN1 gene. In its phosphorylated form, Synapsin I may also be referred to as phosphosynapsin I. Synapsin I is the first of the proteins in the synapsin family of phosphoproteins in the synaptic vesicles present in the central and peripheral nervous systems. Synapsin Ia and Ib are close in length and almost the same in make up, however, Synapsin Ib stops short of the last segment of the C-terminal in the amino acid sequence found in Synapsin Ia. Synapsin II is the collective name for synapsin IIa and synapsin IIb, two nearly identical phosphoproteins in the synapsin family that in humans are encoded by the SYN2 gene. Synapsins associate as endogenous substrates to the surface of synaptic vesicles and act as key modulators in neurotransmitter release across the presynaptic membrane of axonal neurons in the nervous system.

Immunogen: Recombinant protein within human Synapsin I aa 1-705.

Positive control: Mouse brain tissue lysate, Rat brain tissue lysate, mouse primary neuronal, human brain tissue, mouse brain tissue, mouse retina tissue, rat brain tissue, rat retina tissue, IMR-32 cell lysate, U-87 MG cell lysate, Neuro-2a cell lysate.

Subcellular location: Synapse, Golgi apparatus, Presynapse, Cytoplasmic vesicle, secretory vesicle, synaptic vesicle.

Database links: SwissProt: P17600 Human | Q92777 Human | O88935 Mouse | Q64332 Mouse | P09951 Rat | Q63537 Rat

Recommended Dilutions:

WB	1:5,000
IF-Cell	1:500
IHC-Fr	1:500
IHC-P	1:200-1:1,000
IP	1-2µg/sample
IF-Tissue	1:200-1:500

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of Synapsin I + II on different lysates with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/5,000 dilution.

Lane 1: Mouse brain tissue lysate
 Lane 2: Mouse lung tissue lysate (negative)
 Lane 3: Rat brain tissue lysate
 Lane 4: Rat lung tissue lysate (negative)

Lysates/proteins at 20 µg/Lane.


Predicted band size: 74 kDa
 Observed band size: 50-74 kDa

Exposure time: 6 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (HA751346) at 1/5,000 dilution was used in 5% NFDN/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: Immunocytochemistry analysis of mouse primary neuronal cells labeling Synapsin I + II with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/500 dilution.



Cells were fixed in 4% paraformaldehyde for 15 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 15 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/500 dilution in 1% BSA in PBST overnight at 4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 488, HA1121) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (HA601187, red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594, HA1126) was used as the secondary antibody at 1/1,000 dilution.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn




Fig3: Application: IHC-Fr

Species: Mouse

Site: Cerebellum

Sample: Frozen section

Antibody concentration: 1:500

Antigen retrieval: The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for about 2 minutes in microwave oven.




Fig4: Application: IHC-Fr

Species: Mouse

Site: Cerebral cortex

Sample: Frozen section

Antibody concentration: 1:500

Antigen retrieval: The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for about 2 minutes in microwave oven.




Fig5: Application: IHC-Fr

Species: Rat

Site: Cerebral cortex

Sample: Frozen section

Antibody concentration: 1:500

Antigen retrieval: The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for about 2 minutes in microwave oven.




Fig6: Immunohistochemical analysis of paraffin-embedded human brain tissue with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA751346) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig7: Immunohistochemical analysis of paraffin-embedded human lung tissue (negative) with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA751346) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig8: Immunohistochemical analysis of paraffin-embedded mouse brain tissue with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA751346) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig9: Immunohistochemical analysis of paraffin-embedded mouse retina tissue with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA751346) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig10: Immunohistochemical analysis of paraffin-embedded rat brain tissue with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA751346) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig11: Immunohistochemical analysis of paraffin-embedded rat retina tissue with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (HA751346) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.




Fig12: Synapsin I + II was immunoprecipitated from 0.2 mg mouse brain tissue lysate with HA751346 at 2 $\mu\text{g}/10 \mu\text{l}$ beads. Western blot was performed from the immunoprecipitate using HA751346 at 1/1,000 dilution. Mouse Anti-Rabbit IgG kappa light chain secondary antibody (M1208-2) at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: mouse brain tissue lysate (input)

Lane 2: HA751346 IP in mouse brain tissue lysate

Lane 3: Rabbit IgG instead of HA751346 in mouse brain tissue lysate

Blocking/Dilution buffer: 5% NFDm/TBST


Exposure time: 3 seconds; ECL: K1801

Fig13: Western blot analysis of Synapsin I + II on different lysates with Rabbit anti-Synapsin I + II antibody (HA751346) at 1/5,000 dilution.

Lane 1: IMR-32 cell lysate

Lane 2: U-87 MG cell lysate

Lane 3: Neuro-2a cell lysate



Lysates/proteins at 20 $\mu\text{g}/\text{Lane}$.

Predicted band size: 74 kDa

Observed band size: 74/55 kDa

Exposure time: 1 minute 50 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDm/TBST for 1 hour at room temperature. The primary antibody (HA751346) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4 $^{\circ}\text{C}$ overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Forte N et al. Synapsin I Synchronizes GABA Release in Distinct Interneuron Subpopulations. *Cereb Cortex*. 2020 Mar
2. Schwark R et al. Synapsin II Directly Suppresses Epileptic Seizures In Vivo. *Brain Sci*. 2022 Feb

Hangzhou Huan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn