

Anti-IRE1 Antibody [PSH10-60] - BSA and Azide free

HA751363



Product Type: Recombinant Rabbit monoclonal IgG, primary antibodies

Species reactivity: Human, Mouse, Rat, Monkey

Applications: WB

Molecular Wt: Predicted band size: 110 kDa

Clone number: PSH10-60

Description: The serine/threonine-protein kinase/endoribonuclease inositol-requiring enzyme 1 α (IRE1 α) is an enzyme that in humans is encoded by the ERN1 gene. The protein encoded by this gene is the ER to nucleus signalling 1 protein, a human homologue of the yeast Ire1 gene product. This protein possesses intrinsic kinase activity and an endoribonuclease activity and it is important in altering gene expression as a response to endoplasmic reticulum-based stress signals (mainly the unfolded protein response). Two alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Immunogen: Recombinant protein within human IRE1 aa 1-400.

Positive control: HEK-293 cell lysate, Neuro-2a cell lysate, C6 cell lysate, COS-1 cell lysate.

Subcellular location: Endoplasmic reticulum membrane; nucleus.

Database links: SwissProt: O75460 Human | Q9EQY0 Mouse
Entrez Gene: 498013 Rat

Recommended Dilutions:

WB 1:5,000

Storage Buffer: PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co.,Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of IRE1 on different lysates with Rabbit anti-IRE1 antibody (HA751363) at 1/5,000 dilution.



Lane 1: HEK-293 cell lysate
 Lane 2: Neuro-2a cell lysate
 Lane 3: C6 cell lysate
 Lane 4: COS-1 cell lysate

Lysates/proteins at 15 µg/Lane.

Predicted band size: 110 kDa
 Observed band size: 110 kDa

Exposure time: 1 minute; ECL: K1801;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA751363) at 1/5,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Huang R et al. IRE1 signaling regulates chondrocyte apoptosis and death fate in the osteoarthritis. *J Cell Physiol.* 2022 Jan
2. Grandjean JMD et al. Pharmacologic IRE1/XBP1s activation confers targeted ER proteostasis reprogramming. *Nat Chem Biol.* 2020 Oct

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn