

Anti-Cx40 / GJA5 Antibody [PSH10-82] - BSA and Azide free

HA751374



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human, Mouse, Rat, Monkey
Applications:	WB, IP
Molecular Wt:	Predicted band size: 40 kDa
Clone number:	PSH10-82

Description: Gap junction alpha-5 protein (GJA5), also known as connexin 40 (Cx40) — is a protein that in humans is encoded by the GJA5 gene. This gene is a member of the connexin gene family. The encoded protein is a component of gap junctions, which are composed of arrays of intercellular channels that provide a route for the diffusion of low molecular weight materials from cell to cell. Mutations in this gene may be associated with atrial fibrillation. Alternatively spliced transcript variants encoding the same isoform have been described. GJA5 has been identified as the gene that is responsible for the phenotypes observed with congenital heart diseases on the 1q21.1 location. In case of a duplication of GJA5 tetralogy of Fallot is more common. In case of a deletion other congenital heart diseases than tetralogy of Fallot are more common.

Immunogen: Recombinant protein within human GJA5 aa 227-358.

Positive control: HEK-293 cell lysate, HeLa cell lysate, NIH/3T3 cell lysate, F9 cell lysate, COS-1 cell lysate, Mouse lung tissue lysate, Rat lung tissue lysate.

Subcellular location: Cell membrane, Cell junction, gap junction.

Database links: SwissProt: P36382 Human | Q01231 Mouse | P28234 Rat

Recommended Dilutions:

WB	1:2,000
IP	1-2µg/sample

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Images




Fig1: Western blot analysis of Cx40 / GJA5 on different lysates with Rabbit anti-Cx40 / GJA5 antibody (HA751374) at 1/2,000 dilution.

Lane 1: HEK-293 cell lysate (20 µg/Lane)
 Lane 2: HeLa cell lysate (20 µg/Lane)
 Lane 3: NIH/3T3 cell lysate (20 µg/Lane)
 Lane 4: F9 cell lysate (20 µg/Lane)
 Lane 5: COS-1 cell lysate (20 µg/Lane)
 Lane 6: Mouse lung tissue lysate (40 µg/Lane)
 Lane 7: Rat lung tissue lysate (40 µg/Lane)

Predicted band size: 40 kDa

Observed band size: 40 kDa

Exposure time: 25 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA751374) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.




Fig2: Cx40 / GJA5 was immunoprecipitated from 0.2 mg HepG2 cell lysate with HA751374 at 2 µg/10 µl beads. Western blot was performed from the immunoprecipitate using HA751374 at 1/1,000 dilution. HRP Conjugated Anti-Rabbit IgG for IP Nano-secondary antibody at 1/5,000 dilution was used for 1 hour at room temperature.

Lane 1: HepG2 cell lysate (input)
 Lane 2: HA751374 IP in HepG2 cell lysate
 Lane 3: Rabbit IgG instead of HA751374 in HepG2 cell lysate

Blocking/Dilution buffer: 5% NFDM/TBST

Exposure time: 3 seconds; ECL: K1801

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Wang J et al. A novel GJA5 variant associated with increased risk of essential hypertension. Am J Transl Res. 2023 Feb
2. Li J et al. Cis-acting mutation affecting GJA5 transcription is underlying the Melanotic within-feather pigmentation pattern in chickens. Proc Natl Acad Sci U S A. 2021 Oct

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn