



<b>Product Type:</b>	Recombinant Rabbit monoclonal IgG, primary antibodies
<b>Species reactivity:</b>	Human, Mouse, Rat
<b>Applications:</b>	WB
<b>Molecular Wt:</b>	Predicted band size: 74 kDa
<b>Clone number:</b>	PSH11-48

**Description:** Prelamin-A/C, or lamin A/C is a protein that in humans is encoded by the LMNA gene. Lamin A/C belongs to the lamin family of proteins. DNA double-strand damages can be repaired by either homologous recombination (HR) or non-homologous end joining (NHEJ). LMNA promotes genetic stability by maintaining the levels of proteins that have key roles in HR and NHEJ. Mouse cells that are deficient for maturation of prelamin A have increased DNA damage and chromosome aberrations, and show increased sensitivity to DNA damaging agents. In progeria, the inadequacy of DNA repair, due to defective LMNA, may cause features of premature aging (see DNA damage theory of aging).

**Immunogen:** Synthetic phospho-peptide corresponding to residues surrounding Ser22 of Human Lamin A + Lamin C.

**Positive control:** HeLa cell lysate, HeLa treated with 100nM paclitaxel for 20 hours cell lysate, NIH/3T3 cell lysate, NIH/3T3 treated with 100nM paclitaxel for 20 hours cell lysate, C6 cell lysate, C6 treated with 4mM hydroxyurea for 20 hours cell lysate.

**Subcellular location:** Nucleus lamina, Nucleus envelope, nucleoplasm, Nucleus matrix; Nucleus speckle.

**Database links:** SwissProt: P02545 Human | P48678 Mouse | P48679 Rat

**Recommended Dilutions:**  
**WB** 1:2,000


**Storage Buffer:** PBS (pH7.4).

**Storage Instruction:** Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

**Purity:** Protein A affinity purified.

Images

**Fig1:** Western blot analysis of Phospho-Lamin A + Lamin C (S22) on different lysates with Rabbit anti-Phospho-Lamin A + Lamin C (S22) antibody (HA751410) at 1/2,000 dilution and pan Lamin A + Lamin C antibody (ET7110-12) at 1/2,000 dilution.



Lane 1: HeLa cell lysate  
Lane 2: HeLa treated with 100nM paclitaxel for 20 hours cell lysate  
Lane 3: NIH/3T3 cell lysate  
Lane 4: NIH/3T3 treated with 100nM paclitaxel for 20 hours cell lysate  
Lane 5: C6 cell lysate  
Lane 6: C6 treated with 4mM hydroxyurea for 20 hours cell lysate  
Lane 7: HeLa treated with 100nM paclitaxel for 20 hours cell lysate, then the membrane treated with λpp for 1 hour  
Lane 8: NIH/3T3 treated with 100nM paclitaxel for 20 hours cell lysate, then the membrane treated with λpp for 1 hour  
Lane 9: C6 treated with 4mM hydroxyurea for 20 hours cell lysate, then the membrane treated with λpp for 1 hour

Lysates/proteins at 20 µg/Lane.

Predicted band size: 74 kDa  
Observed band size: 70/65 kDa

Exposure time: 4 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA751410) at 1/2,000 dilution and pan Lamin A + Lamin C antibody (ET7110-12) at 1/2,000 dilution were used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

**Note:** All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Background References

1. Kovacs MT et. al. DNA damage induces nuclear envelope rupture through ATR-mediated phosphorylation of lamin A/C. Mol Cell. 2023 Oct
2. Yamada S et al. TEAD1 trapping by the Q353R-Lamin A/C causes dilated cardiomyopathy. Sci Adv. 2023 Apr

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物  
HUABIO  
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation