

Anti-BATF3 Antibody [PSH13-52] - BSA and Azide free HA751471



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Human
Applications:	WB, IP, ChIP
Molecular Wt:	Predicted band size: 14 kDa
Clone number:	PSH13-52

Description: AP-1 family transcription factor that controls the differentiation of CD8+ thymic conventional dendritic cells in the immune system. Required for development of CD8-alpha+ classical dendritic cells (cDCs) and related CD103+ dendritic cells that cross-present antigens to CD8 T-cells and produce interleukin-12 (IL12) in response to pathogens (By similarity). Acts via the formation of a heterodimer with JUN family proteins that recognizes and binds DNA sequence 5'-TGA[CG]TCA-3' and regulates expression of target genes.

Immunogen: Recombinant protein within human BATF3 aa 1-127.

Positive control: U-937 cell lysate, HDLM2 cell lysate.

Subcellular location: Nucleus.

Database links: SwissProt: Q9NR55 Human

Recommended Dilutions:

WB	1:5,000
IP	1-2µg/sample
ChIP	Use 0.5~2 µg for 25 µg of chromatin.

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of BATF3 on different lysates with Rabbit anti-BATF3 antibody (HA751471) at 1/5,000 dilution.



Lane 1: U-937 cell lysate
 Lane 2: Ramos cell lysate (negative)
 Lane 3: HDLM2 cell lysate
 Lane 4: HepG2 cell lysate (negative)


Lysates/proteins at 20 µg/Lane.

Predicted band size: 14 kDa
 Observed band size: 18 kDa

Exposure time: 3 minutes; ECL: K1801;
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA751471) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Fig2: BATF3 was immunoprecipitated from 0.2 mg U-937 cell lysate with HA751471 at 2 µg/10 µl beads. Western blot was performed from the immunoprecipitate using HA751471 at 1/5,000 dilution. HRP Conjugated Anti-Rabbit IgG for IP Nano-secondary antibody at 1/5,000 dilution was used for 1 hour at room temperature.



Lane 1: U-937 cell lysate (input)
 Lane 2: HA751471 IP in U-937 cell lysate
 Lane 3: Rabbit IgG instead of HA751471 in U-937 cell lysate

Blocking/Dilution buffer: primary antibody dilution (K1803)
 Exposure time: 3 minutes; ECL: K1801




Fig3: Chromatin immunoprecipitations were performed with cross-linked chromatin from U-937 cells with BATF3 (HA751471) or Normal Rabbit IgG according to the ChIP protocol. The enriched DNA was quantified by real-time PCR using indicated primers. The amount of immunoprecipitated DNA in each sample is represented as signal relative to the total amount of input chromatin, which is equivalent to one.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
 HUABIO
 www.huabio.cn

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Jain N et al. TET2 guards against unchecked BATF3-induced CAR T cell expansion. Nature. 2023 Mar
2. Ziblat A et al. Batf3(+) DCs and the 4-1BB/4-1BBL axis are required at the effector phase in the tumor microenvironment for PD-1/PD-L1 blockade efficacy. Cell Rep. 2024 May

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUAABIO
www.huabio.cn