

Anti-Phospho-AMPK alpha 1 (T183) +AMPK alpha 2 (T172) Antibody [PSH20-31] - BSA and Azide free

HA751747



Product Type: Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity: Human
Applications: WB
Molecular Wt: Predicted band size: 64 kDa
Clone number: PSH20-31

Description: Catalytic subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators

Positive control: HEK-293 treated with 200nM Calyculin A for 1 hour cell lysate.

Subcellular location: Cytoplasm, Nucleus.

Database links: SwissProt: Q13131 Human | P54646 Human

Recommended Dilutions:
WB 1:2,000

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation


Images

Fig1: Western blot analysis of Phospho-AMPK alpha 1 (T183) +AMPK alpha 2 (T172) on different lysates with Rabbit anti-Phospho-AMPK alpha 1 (T183) +AMPK alpha 2 (T172) antibody (HA751747) at 1/2,000 dilution.

Lane 1: HEK-293 cell lysate

Lane 2: HEK-293 treated with 200nM Calyculin A for 1 hour cell lysate

Lane 3: HEK-293 treated with 200nM Calyculin A for 1 hour cell lysate, then the membrane treated with λ pp for 1 hour



Lysates/proteins at 20 μ g/Lane.

Predicted band size: 64 kDa

Observed band size: 64 kDa

Exposure time: 2 minutes; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDN/TBST for 1 hour at room temperature. The primary antibody (HA751747) at 1/2,000 dilution was used in primary antibody dilution (K1803) at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

- Albayrak G et al. Memantine shifts cancer cell metabolism via AMPK1/2 mediated energetic switch in A549 lung cancer cells. EXCLI J. 2021 Feb
- Lin FC et al. Protective Effects of Kirenol against Lipopolysaccharide-Induced Acute Lung Injury through the Modulation of the Proinflammatory NFkappaB Pathway and the AMPK2-/Nrf2-Mediated HO-1/AOE Pathway. Antioxidants (Basel). 2021 Jan

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn