

Anti-CD38 Antibody [PSH20-33] - BSA and Azide free

HA751749



Product Type:	Recombinant Rabbit monoclonal IgG, primary antibodies
Species reactivity:	Mouse
Applications:	WB
Molecular Wt:	Predicted band size: 34 kDa
Clone number:	PSH20-33

Description: CD38 is a type II integral membrane glycoprotein which is present on early B and T cell lineages and activated B and T cells but is absent from most mature resting peripheral lymphocytes. CD38 is also found on thymocytes, pre-B cells, germinal center B cells, mitogen-activated T cells, monocytes and Ig-secreting plasma cells. CD38 acts as a NAD glycohydrolase in T lymphocytes. On hematopoietic cells CD38 induces activation, proliferation, and differentiation of mature T and B cells and mediates apoptosis of myeloid and lymphoid progenitor cells. In addition to acting as a signaling receptor, CD38 is also an enzyme capable of producing several calcium-mobilizing metabolites, including cyclic adenosine diphosphate ribose (cADPR). CD38 also plays a role in maintaining survival of an invariant NK T (iNKT) cell subset that preferentially contributes to the maintenance of immunological tolerance.

Positive control: Mouse spleen tissue lysate, Mouse small intestine tissue lysate.

Subcellular location: Cell surface, Membrane.

Database links: SwissProt: P56528 Mouse

Recommended Dilutions:

WB 1:5,000

Storage Buffer: 1*PBS (pH7.4).

Storage Instruction: Store at +4℃ after thawing. Aliquot store at -20℃. Avoid repeated freeze / thaw cycles.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

Images

Fig1: Western blot analysis of CD38 on different lysates with Rabbit anti-CD38 antibody (HA751749) at 1/5,000 dilution.

Lane 1: Mouse spleen tissue lysate
Lane 2: Mouse small intestine tissue lysate


Lysates/proteins at 40 µg/Lane.

Predicted band size: 34 kDa
Observed band size: 45 kDa

Exposure time: 10 seconds; ECL: K1801;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (HA751749) at 1/5,000 dilution was used in primary antibody dilution (K1803) at 4℃ overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody (HA1001) at 1/50,000 dilution was used for 1 hour at room temperature.



Note: All products are “FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE”.

Background References

1. Yang Q et al. NADase CD38 is a key determinant of ovarian aging. Nat Aging. 2024 Jan
2. Yu S et al. CD38-Targeting Peptide Vaccine Ameliorates Aging-Associated Phenotypes in Mice. Aging Cell. 2025 Sep

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation