

Anti-Apolipoprotein A1 Antibody

M0809-13



Product Type:	Mouse monoclonal IgG1, primary antibodies
Species reactivity:	Human
Applications:	WB, IF-Cell, IHC-P
Molecular Wt:	Predicted band size: 31 kDa

Description: Apolipoprotein AI (Apo-AI) is a protein that in humans is encoded by the APOA1 gene. As the major component of high-density lipoprotein (HDL) particles, it has a specific role in lipid metabolism. Apolipoprotein AI is the major protein component of high density lipoprotein (HDL) particles in plasma. Chylomicrons secreted from the intestinal enterocyte also contain Apo-AI, but it is quickly transferred to HDL in the bloodstream. The protein, as a component of HDL particles, enables efflux of fat molecules by accepting fats from within cells (including macrophages within the walls of arteries which have become overloaded with ingested fats from oxidized low-density lipoprotein (LDL) particles) for transport (in the water outside cells) elsewhere, including back to LDL particles or to the liver for excretion.

Immunogen: Synthetic peptide within human Apolipoprotein A1 aa 19-68.

Positive control: Human plasma lysates, human liver tissue.

Subcellular location: Secreted.

Database links: SwissProt: P02647 Human

Recommended Dilutions:

WB	1:500-1:1,000
IF-Cell	1:50-1:200
IHC-P	1:50-1:200

Storage Buffer: 1*PBS (pH7.4), 0.2% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.

Storage Instruction: Store at +4°C.

Purity: Protein A affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880


Technical:0086-571-89986345

Service mail:support@huabio.cn

 华安生物
HUABIO
www.huabio.cn

Images

Fig1: Western blot analysis of Apolipoprotein A1 on human plasma lysates with Mouse anti-Apolipoprotein A1 antibody (M0809-13) at 1/1,000 dilution.



Lysates/proteins at 20 µg/Lane.

Predicted band size: 31 kDa

Observed band size: 26 kDa

Exposure time: 5 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (M0809-13) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Mouse IgG - HRP Secondary Antibody (HA1006) at 1/50,000 dilution was used for 1 hour at room temperature.




Fig2: Immunohistochemical analysis of paraffin-embedded human liver tissue with Mouse anti-Apolipoprotein A1 antibody (M0809-13) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (M0809-13) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Background References

1. Xu X et al. Apolipoprotein A1-Related Proteins and Reverse Cholesterol Transport in Antiatherosclerosis Therapy: Recent Progress and Future Perspectives. *Cardiovasc Ther.* 2022 Jan
2. Maarfi F et al. A Study on Multiple Facets of Apolipoprotein A1 Milano. *Appl Biochem Biotechnol.* 2023 Jul

Hangzhou Huaan Biotechnology Co., Ltd.

Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

华安生物
HUABIO
www.huabio.cn