Anti-S tag Antibody

R1401-17



Product Type:	Rabbit polyclonal IgG, primary antibodies
Species reactivity:	Species independent
Applications:	WB, ELISA
Description:	S-tag is the name of an oligopeptide derived from pancreatic ribonuclease A. The amino acid sequence of the S-tag is: Lys-Glu-Thr-Ala-Ala-Ala-Lys-Phe-Glu-Arg-Gln-His-Met-Asp-Ser. It is believed that the peptide with its abundance of charged and polar residues could improve solubility of proteins it is attached to. This antibody recognizes the s-tag fused to either the amino- or carboxy-terminus of targeted proteins.
lmmunogen:	Recombinant protein fused with S Tag.
Positive control:	S tagged fusion protein
Recommended Dilutions: WB ELISA	1:5,000-1:10,000 1:10,000
Storage Buffer:	1*PBS (pH7.4), 0.2% BSA, 40% Glycerol. Preservative: 0.05% Sodium Azide.
Storage Instruction:	Store at +4 $^\circ\!\!C$ after thawing. Aliquot store at -20 $^\circ\!\!C$ or -80 $^\circ\!\!C$. Avoid repeated freeze / thaw cycles.
Purity:	Immunogen affinity purified.

Hangzhou Huaan Biotechnology Co., Ltd.




S345 Service mail:support@huabio.cn



Orders:0086-571-88062880 Techr

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation

R1401-17 - Page 2



Note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE".

Hangzhou Huaan Biotechnology Co., Ltd.



Orders:0086-571-88062880

Technical:0086-571-89986345

Service mail:support@huabio.cn

Applications:WB=Western blot IHC-P=Immunohistochemistry (paraffin) IF-Cell=Immunofluorescence (Cell) IF-Tissue=Immunofluorescence (Tissue) FC=Flow cytometry IP=Immunoprecipitation